Assessment and Analysis of two Pinus kesiya Provenance Trials in Indonesia Trial No. 7 (T 72) Aek Nauli, Trial No. 8 (T 70) Habinsaran

Hansen, Christian Pilegaard; Hansen, Christian Pilegaard; Ræbild, Anders; Saragih, Thomas

Publication date: 2003

Document version Publisher's PDF, also known as Version of record

Citation for published version (APA): Hansen, C. P., Hansen, C. P., Ræbild, A., & Saragih, T. (2003). Assessment and Analysis of two Pinus kesiya Provenance Trials in Indonesia: Trial No. 7 (T 72) Aek Nauli, Trial No. 8 (T 70) Habinsaran. Danida Forest Seed Centre. Results and Documentation No. 21

Assessment and analysis of two *Pinus kesiya* provenance trials in Indonesia

Trial No. 7 (T 72) Aek Nauli Trial No. 8 (T 70) Habinsaran

by

Christian Pilegaard Hansen², Anders Ræbild² and Thomas Saragih¹

Research & Development Department, RGMI Forestry¹, North Sumatra, Indonesia Danida Forest Seed Centre², Denmark



Results and Documentation No. 21

Danida Forest Seed Centre

August 2003

Citation:

Hansen, C.P., A. Ræbild and T. Saragih. 2003. Assessent and analysis of two *Pinus kesiya* trials in Indonesia. Results and Documentation No. 21. Danida Forest Seed Centre, Humlebaek, Denmark.

Reproduction is allowed with citation

ISSN 0902-3224

Cover photo:

Poor branching characteristics; whorls of heavy persistent branches and long internodes have militated against the acceptance of *P. kesiya* as a plantation species. Here Habinsaran trial site, Indonesia. Photo: Christian Pilegaard Hansen

This publication can be requested from:

Danida Forest Seed Centre Krogerupvej 21. DK-3050 Humlebaek, Denmark Phone: +45-49190500 Fax: +45-49190258 Email: dfsc@sns.dk Web Site: www.dfsc.dk

and/or be downloaded from the DFSC homepage: www.dfsc.dk/publications/

Technical Editor: Melita Jørgensen

Print: Toptryk A/S, Graasten

Results and documentations are publications of analyses of e.g. provenance trials, carried out between DFSC and other institutions. DFSC publications are distributed free of charge

Danida Forest Seed Centre (DFSC) is a Danish non-profit institute which has been working with development and transfer of know-how in management of tree genetic resources since 1969. The development objective of DFSC is to contribute to improve the benefits of growing trees for the well-being of people in developing countries. DFSC's programme is financed by the Danish International Development Assistance (Danida).

Preface

This report presents the results of a joint assessment of two *Pinus kesiya* provenance trials in Indonesia. The trials were established by RGMI Forestry, Research and Development Division in 1992 as part of an international series of provenance trials of the species.

The joint RGMI/DFSC field assessment took place in September 1999 with participation of Thomas Saragih, Wagiman, Nabil, Parlindungan Panjaitan and Gibson Manurung of RGMI Forestry, R&D. From DFSC participated Anders Ræbild and Christian Pilegaard Hansen.

Paul Clegg, Dr. Mok Chak Kim and Dr. Chan Yik Kuan of RGMI, R&D are thanked for their kind assistance in planning and arrangements for the field assessment.

Useful comments and assistance in the statistical analysis and interpretation of results were received from Anders Ræbild and Erik D. Kjaer, DFSC.

Contents

Prefac Cont Acror	ce ents nyms and abbreviations	i iii iv
1	Background	1
2	P. kesiya provenance trials established by RGMI Forestry, Indonesia	2
3	Field assessment and data management	4
4	Statistical analysis	5
	4.1 Plot of raw data	6
	4.2 Statistical model	6
	4.3 Co-variates	6
	4.4 Check of model assumptions	7
	4.5 What to do when model assumptions are not fulfilled	8
	4.6 Fixed or random effects	9
	4.7 Test of differences between species and provenances	9
	4.8 Lsmeans (estimates from the fixed model)	10
	4.9 Best Linear Unbiased Predictors (BLUPs - estimates from the random model)	10
5	Results of statistical analysis of individual traits	11
	5.1 Survival	12
	5.2 Height growth	14
	5.3 Diameter growth	16
	5.4 Mean volume of tree	18
	5.5 Total volume per hectare	20
	5.6 Stemform	22
	5.7 Wood density (Pilodyn)	24
	5.8 Branching (branch diameter)	26
	5.9 Foxtailing	28
	5.10 Flowering	30
6	Conclusions	32
	6.1 Growth	32
	6.2 Adaptation	32
	6.3 Quality	32
	6.4 Overall conclusion	32
7	References	34

Annexes

Annex 1MapsAnnex 2Trial descriptionsAnnex 3Site descriptionsAnnex 4Plot data set

Acronyms

CAMCORE	Central America and Mexico Coniferous Resources Co-operative, USA
CIEF	Centro de Investigaciones y Experiencias Forestales, Argentina
CSIRO	Commonwealth Scientific and Industrial Research Organisation, Australia
Danida	Danish International Development Assistance
DFSC	Danida Forest Seed Centre, Humlebæk, Denmark
EMBRAPA	Empresa Brasileira de Pesquisa Agropecuária, Petrolona, Pernambuca, Brazil
FAO	Food and Agriculture Organization of the United Nations, Rome, Italy
ICFRE	Indian Council of Forestry Research and Education, Dehra Dun, India
IF	Instituto Florestal, São Paulo, Brazil
IPEF	Instituto de Pesquiasa e Estudos Florestais, Piracicaba, Brazil
OFI	Oxford Forestry Institute, United Kingdom
RCFTI	Research Centre for Forest Tree Improvement, Forest Science Institute, Vietnam
RFD	Royal Forest Department, Thailand

1. Background

The Aek Nauli and Habinsaran trials form part of an international series of provenance trials of *P. kesiya*.

The objective of the international series is to explore and analyse the genetic variation in growth, quality and adaptive traits among provenances of *P. kesiya* throughout the range of the species. The results will facilitate an informed choice of seed source in planting programmes. Furthermore, the results will be useful when planning conservation activities of the species.

Below the background of the international series is briefly described.

Initial research on inter-population differences in *P. kesiya* was undertaken in Zambia in the 1950s. The test material included provenances from the Philippines, Vietnam and Assam. A comprehensive review of these studies is given in Armitage and Burley (1980).

During 1969, FAO and the Forest Research Institute of Australia sponsored seed collections of 19 seed sources of P. kesiya from the Philippines (17 provenance collections and 2 commercial seedlots). The material was complemented by two Zambian land races (of Philippine and Vietnamese origin, respectively). These collections were used for provenance trials in a large number of countries for which the Commonwealth Forestry Institute supplied advice and assisted in data processing and interpretation (Burley and Wood 1976). Results from individual trials were reviewed by Gibson and Barnes (1984). They concluded that neither provenance representation, nor test site representation warranted an international evaluation. It was recommended that a more comprehensive exploration and analysis of the genetic variation of P. kesiya should be undertaken. Recommendations in this regard was also put forward by the Sixth Session of the FAO Panel of Experts on Forest Gene Resources (FAO 1988).

Exploration of provenance variation and collection of seed for field trials took place in the late 1980s in collaboration between national institutions in Brazil, Myanmar, China, Madagascar, Philippines, Thailand, Vietnam, Zambia, Zimbabwe, Oxford Forestry Institute (OFI) and Danida Forest Seed Centre (DFSC). In 1988, seed collections were complete and distribution to collaborating countries could begin (Barnes et al. 1989). Distribution of seed was co-ordinated by OFI and handled by DFSC. During 1989-93, seed of 42 provenances and land races from the above 9 countries were distributed to 20 institutions in 19 countries. Some of the seedlots were separated by mother trees to allow testing of individual families.

A status of seed distribution and established field trials is found in DFSC (1996) and DFSC (1997). Some 30 trials have been established in 17 countries. Trials in Argentina, Brazil, Colombia, Indonesia, South Africa, Swaziland, Vietnam and Zimbabwe are reported with high survival and are in general in good conditions. Status of trials in Burundi, India, Rwanda and Sri Lanka is unknown, as no information has been received from these countries. Trials established in Fiji, Kenya, Nepal, the Philippines and Thailand have been abandoned because of fire damage, drought and browsing.

In a circular letter sent out by OFI and DFSC in 1996, host institutes were asked about their interest in undertaking a joint evaluation and were at the same time asked about the status of the trials (DFSC 1996). Positive responses in regard to the proposal of undertaking a joint assessment and analysis of trials have been received from all countries where existence of trials has been confirmed. The number, distribution and representation of provenances in these trials were considered sufficient to justify an assessment and analysis of the international series. Of special interest is the possibility of an in-depth analysis of provenance x site interactions, thanks to the representation of the same set of provenances at many trial sites.

A manual was elaborated during 1997-98 with a proposal for a set of characters to be assessed in all trials (DFSC 1998). Field assessment of trials commenced in April 1998 (Vietnam).

2. *P. kesiya* provenance trials in Indonesia

RGMI Forestry, Indonesia, has established two provenance trials of *P. kesiya* in 1992 in North Sumatra, Indonesia. The trials are located at the Aek Nauli and Habinsaran forestry sectors and have trial identification numbers T72 and T70, respectively.

The provenances represented in the trials are shown in the below table.

Remarks about the table:

- 1. Seedlots 712, 713, 714, 715, 716, 183, 118, 366, i.e. the not *P. kesiya/P. yunnanensis* sources, are local controls, and do not form part of the international seed exchange under the international programme.
- 2. The field assessment revealed that the Doi Inthanon provenance of *P. kesiya* in fact was *P. tecunumanii*. As the seed lot could not be further identified, it has been omitted from the analysis.
- 3. The Guanaja provenance of *P. caribaea* var. *hondurensis* and the Coto Mines provenance of *P. kesiya* are only in the Aek Nauli trial, not in the Habinsaran one.
- 4. Eucalyptus grandis has been included as a local control in both trials. The Eucalyptus plots in the Aek Nauli trial have been cut down, and hence Eucalyptus is only in the Habinsaran trial. The plots have been assessed, but not included in the statistical analysis.

5. The Simao provenance of *P. kesiya* is only present in the Habinsaran trial, and only in one of the four replicates. This plot has been assessed, but the provenance has not been included in the statistical analysis.

Details of the trial establishment and management are presented in Annex 2 and descriptions of the sites are in Annex 3.

The Aek Nauli trial has a low survival rate. Many trees have reportedly been cut down during weeding operations, probably because weedings have been delayed. It is further reported that trees in the trial have suffered from herbicide application. As a result, many plots in the Aek Nauli trial have few trees left, and other plots have been entirely lost (no trees left). This is weakening the statistical analysis of the trial.

The Habinsaran trial, on the other hand, has a much higher and more uniform survival. The growth at the Habinsaran trial also compares favourable to the Aek Nauli trial.

As mentioned above, both trials suffer from misplaced seedling/rows/plots. The problem with seedlot 723 (the Doi Inthanon source of *P. kesiya*) in both trials has been mentioned above, but also in other plots, there are rows, parts of rows or single trees of other origin, i.e. misplaced seedlings. These trees have in all cases been omitted from the assessment.

Local ID	DFSC Acc. No.	Species	Provenance	Country
712	-	P. oocarpa	Mal Paso	Guatemala
713	-	P. tecunumanii	Mt. Pine Ridge	Belize
714	-	P. oocarpa	El Paraiso	Honduras
715	-	P. tecunumanii	San Raphael	Nicaragua
716	-	P. caribaea	Guanaja	Honduras
718	1572/85	P. kesiya	Coto Mines	Philippines
719	1525/85	P. kesiya	Nam Now	Thailand
720	1521/85	P. kesiya	Nong Krating	Thailand
721	1519/85	P. kesiya	Lang Hanh	Vietnam
722	1522/85	P. kesiya	Doi Suthep	Thailand
723	1523/85	P. kesiya	Doi Inthanon	Thailand
724	1639/86	P. kesiya	Simao	China
725	1783/88	P. kesiya	Bodana A8	Madagascar
726	1773/88	P. kesiya	Aungban	Myanmar
727	1633/86	P. yunnanensis	Shangsi	China
183	-	P. merkusii	Indonesia	Indonesia
118	-	P. patula	Zimbabwe landrace	Zimbabwe
366	-	E. grandis	Coff's Harbour	Australia

3. Field assessment and data management

The assessment followed the methodology described in DFSC (1998) and included the characters:

- 1. Survival;
- 2. Health;
- 3. Social status (Kraft);
- 4. Height;
- 5. Diameter (DBH);
- 6. Straightness;
- 7. No. of whorls;
- 8. No. of branches in whorl;
- 9. Branch diameter;
- 10. No. of forks;
- 11. Position of first fork;
- 12. Foxtail;
- 13. Flowering and fruiting;
- 14. Wood density (Pilodyn);

For a detailed description of the assessment methodology, please refer to DFSC (1998).

The assessment was a full assessment, i.e. all trees within each plot were included. For the local controls, i.e. seedlots not *P. kesiya*, the assessment was limited to survival, height, diameter, stemform and pilodyn.

Relative wood density was measured with a Pilodyn wood tester with pin diameter 2.0 mm.

Assessment data was recorded in the field on assessment sheets, see example in DFSC (1998). The data was immediately after the assessment entered to a lap-top computer in spreadsheet format. From the spreadsheet, data was later transferred to a SAS-dataset for further analysis.

4. Statistical analysis

Overview of steps involved in the statistical analysis



The objectives of the statistical analysis are:

- to examine statistically significant differences between seedlots (provenances) in adaptability, growth and quality traits. A list of analysed traits is provided in Chapter 5;
- to conclude and recommend on the practical application of the results (species and provenance recommendations);
- to investigate patterns of genetic variation;
- to provide data for an overall analysis of the international series of provenance trials of *Pinus kesiya*, i.e. analysis across sites. This step will await completion of the analysis of individual trials.

Statistical analysis is done on plot values, e.g. plot averages or plot sums. Calculation of plot values is described in Annex 4.

The SAS analytical package has been used for the analysis (SAS, 1990).

The statistical analysis of each trait follows a sequence of steps. They are:

- 1. Plot of raw data;
- 2. Formulation of statistical model;
- 3. Test of co-variates;
- 4. Check of model assumptions;
- 5. When model assumptions are not fulfilled: (a) transformation of data; (b) deletion of outliers; (c) weight statement;
- 6. Test of differences between species and provenances;
- 7. Calculation of lsmeans (estimated from a model with fixed effects);
- 8. Calculation of BLUPs (estimated from a model with random effects).

The statistical analysis is illustrated in the above figure and the steps are further described in the below text.

Generally speaking, two different approaches are applied in the statistical analysis: a fixed effect approach and a random effect approach. The fixed effects approach is concerned with the genetic entries (seedlots) actually in the trial, whereas the random effects approach concerns what would happen if the experiment was to be repeated. Following the fixed effect approach, the estimates are calculated as least square means (lsmeans), whereas the random effect approach gives the best linear unbiased predictors (BLUPs). See further explanation below.

4.1 Plot of raw data

The main purpose of the plots is to indicate the scale of the variable along with a first impression of the variation within the trial. Often the visual inspection of the data reveals clear differences between the provenances, or gives hints regarding proper transformations of the data. Obvious outliers (extreme values) may also be identified already at this stage. The most useful single plot is probably a plot of the variable against the provenances, marking the values with values identifying the blocks. However, other plots may also be relevant, e.g. plotting the values by block or by the distance along the axis of the trial.

4.2 Statistical model

The test of differences between seedlots (provenances) is based on the model:

$$X_{ik} = \mu + provenance_i + block_k + \varepsilon_{ik}$$

where X_{jk} is the value of the trait in question (e.g. height) in plot *jk*,

 $\boldsymbol{\mu}$ is the grand mean,

provenance, is the effect of seedlot number *J* and is assumed to be either a fixed or a random effect, according to which approach is used (see later),

 $block_k$ is the effect of block k in the trial, assumed to be a random effect, and

 ε_{jk} is the residual of plot *jk* and is assumed to follow a normal distribution $N(0, \sigma_{c}^{2})$.

Please note that the controls (seedlots not *Pinus kesiya*/*P.yunnanensis*) are considered (analysed) together with these sources not considering that they actually are sources of different species.

4.3 Co-variates

In order to reduce the residual variation in trials with heterogeneous trial conditions (e.g. variation in soil, elevation, slope and exposure within trial), a number of co-variates are included in the model. As a standard routine the following four co-variates are tested:

- plotx: Horizontal position of plot within trial (see map of trial);
- ploty: Vertical position of plot within trial (see map of trial);

To catch non-linear patterns of site variation vertically and horizontally, plotx2 and ploty2 are applied:

plotx2: plotx2=(plotx - mean(plotx))²
ploty2: ploty2=(ploty - mean(ploty))²

In addition to the above four co-variates, additional co-variates are considered in some of the trials:

level: Level of plot in relation to a reference plot within the trial (0);

plotxy

$Plotxy = plotx \ x \ ploty$

In testing the effect of co-variates, we start with a model with all co-variates included. Co-variates that are not significant are removed successively by removing the least significant co-variate and running the model again until all remaining covariates in the model are significant (P<0.10).

4.4 Check of model assumptions

The statistical model rests on a number of standard assumptions. Key assumptions are (see e.g. Box *et al.* 1978):

- (i) that the residuals are independent;
- (ii) that the residuals follow a normal distribution;
- (iii) that there is variance homogeneity in effects included in the model.

The model assumptions are checked graphically by producing a number of plots:

- 1. Student's residuals versus predicted values;
- 2. Cooks distance versus predicted values;
- 3. Student's residuals versus provenance;
- 4. Frequency chart of residuals;
- 5. Student's residuals versus block;
- 6. Student's residuals versus plotx;
- 7. Student's residuals versus ploty;
- 8. Student's residuals versus level (if level is among the considered co-variates).

The residuals represent variation that can not be accounted for by the model. For each observation, the model calculates a predicted value, taking into account the effects of the model (provenance, block and co-variates). The residual variation is then the difference between the observed value and the predicted value.

Student's residuals (also called 'standardised residuals') are calculated as the residual divided by its standard error. If the assumption of normal distributed residuals is valid, the Student's residuals have the property of a normal distribution with mean 0 and variance 1, meaning that 95% of the values should lie within \pm 1.96. In cases of trials with imbalance, the Student's residuals correct for imprecision due to low sample numbers, and in models with co-variates they compensate for large deviations at extreme values.

The Student's residual e_i for observation ij is given by

$$e_t = \frac{e_{ij}}{s_{ij}} = \frac{X_{ij} - P_{i\bullet} - B_{\bullet j}}{s_{ij}}$$

where e_{ij} is the residual, X_{ij} is the value for observation ij, $P_{i,\bullet}$ is the effect of provenance i, $B_{\bullet,j}$ is the effect of block j, and s_{ij} is the standard deviation (standard error) of observation ij.

Cooks distance gives a measure of the influence of a single observation (plot) on the model, and gives an indication of possible 'outliers' (see below) (Afifi & Clark 1996). A high value indicates an observation with a large influence on the outcome of the model.

In the following, a description of the check of the model assumptions is given.

Independence

The assumption of independence means that the residual of one observation is not dependent on the residual of another. This assumption is typically violated when using pseudo-replicates, e.g. when doing more observations on the same experimental unit and treating them as different experimental units. Another example is when two or more plots of the same provenance within the same block are treated as independent observations. In such cases, an average of the values should be used as the block value for the provenance in question.

The graphical check of the residuals does not reveal possible problems with observations dependent upon each other, and there is no easy method to ensure that the condition of independence is fulfilled. Proper design and planning of the experiment result and application of a correct statistical model is the best insurance to obtain independent observations.

The assumption of independence may also be violated if there is a time- or site-dependency in the data. To check for such dependency, residuals are plotted against the horizontal and vertical axis of the trial (plotx and ploty) and where applicable, also the level of plot, to investigate any systematic environmental variation. Usually there is none, as the co-variates (plotx and ploty) account for this.

Normality

The assumption of normality may be checked in various ways, graphically as well as by statistical tests. In this analysis, we use the frequency chart of residuals as a graphical check. In the frequency chart, the frequencies should be more or less bellshaped with no large tails at the ends. A formal statistical test, the Shapiro-Wilk statistic, is given in the SAS-procedure UNIVARIATE with the option NORMAL (SAS 1988a). This procedure also offers different kinds of plots of the residuals. However, since the test is usually considered to be conservative, rejecting only severe deviations from normality, the test results should be considered with caution (Brockhoff, pers. comm.).

When the number of observations is low, it becomes increasingly difficult to check the assumption of normality. Even though the frequency chart may show a rather odd and irregular distribution, this need not be a sign of non-normality. At small sample sizes it is not unlikely that odd distributions may result from random variation, and unless the test for normality demonstrates that the assumption is violated, there is no need to reject the model. On the other hand, when the number of samples is very large, the test for normality may become rejected even though the frequency chart of residuals appears to be normal. This is because the power of the test increases with the number of observations, and even small deviations from normality may result in rejection of the hypothesis of normal distributed residuals. In such cases it

should be considered whether the frequency chart indicate that the assumption is fulfilled, or the deviations are so large that transformation of data (see later) is required.

Deviations from the assumption of normality may also be interpreted as a distribution with a large number of outliers (see later).

Variance inhomogeneity

Variance inhomogeneity occurs when different experimental units (blocks and provenances) have different variance. A typical example is when the residuals of some provenances appear very clustered in the diagram of Student's residuals versus provenances, whereas the residuals of other provenances are spread out, often with values of Student's residuals exceeding ± 2 . This may result from a simple scale effect (larger provenances have larger variance), in which case the plots of Student's residuals and Cook's distance versus predicted values appear funnel-shaped. It may also be related to the provenance itself (some provenances are more variable than others). In this case, the variance inhomogeneity will be displayed in the plot of student's residuals versus provenance.

Outliers

Outliers are extreme observations that do not follow the trends of the remaining data. Such observations may have a large influence on the estimates and statistical tests of the model and should therefore be considered carefully.

Outliers are detected by inspection of the plots of Student's residuals and of Cook's distance. Observations that have values of Student's residuals exceeding ± 2.5 (rule of thumb), and observations with large values of Cook's distance, are possible outliers and should be investigated further. Outliers may be due to errors in the recording or typing of data, or due to mislabelling of the seedlots in the nursery or in the field. Poor survival in the plot, leaving only a few trees to use in the calculation of plot means is another source of outliers. However, it also happens that the outliers are due to some unexplained variation, perhaps in soil conditions or other environmental variation. Finally it should be mentioned that a large number of outliers might indicate that the distribution of residuals is not normal, and hence that a transformation of data is required (see later).

When outliers occur as a result of errors, the dataset should of course be corrected, which will solve the problem. It is less obvious what to do in the cases where there are no easy explanations. Outliers should only be excluded if it can be justified, i.e. an explanation can be given. In a few cases, however, explanations were not found, and observations were excluded alone on basis of the extreme nature of the value. Great care is required in the decision to exclude plot values, as it will have great importance for the result of the analysis, especially with few blocks (replications). Running the analysis again without the outlier(s) gives an indication of the sensitivity of the analysis in regard to the outliers, and assist in deciding whether to keep or delete the extreme observation(s).

In the interpretation of the statistical analysis in this report, it is always mentioned if one or more extreme values have been considered as outliers and omitted from the analysis, and on what grounds.

4.5 What to do when model assumptions are not fulfilled

In many cases one or more of the model assumptions are not fulfilled. In the below, procedures for correction are described.

Independence

Apart from making sure that the statistical design and the model is correct there is not much to do about dependence between observations. If some clear variation can be observed in the residuals, other co-variates than the ones mentioned above could be considered.

Deviations from normality

Usually deviations from normality are handled by transformation of data. Snedecor & Cochran (1980) and Afifi & Clark (1996) provide guidance on data transformations:

- 1. Counts (of rare events) often follow a Poisson distribution and are transformed with the square root.
- 2. Variables having a binomial character (e.g. dead or alive) summarised in a proportion (e.g. living trees in a plot) may be transformed with the arc sine transformation.
- 3. If the standard deviation varies directly with the mean, a logarithmic transformation may stabilise the variance.

There are theoretical reasons for choosing the above transformations (Snedecor & Cochran 1980), but it follows from Afifi & Clark (1996) that the range of transformations may be seen as a continuum and that various other transformations are available.

None of the variables included in the present assessment have the character of a Poisson distribution, but the square root transformation has nevertheless in some cases been applied.

In many cases the analysis of survival data results in skewed distributions of the residuals, with tails at either the lower or upper end (many trees dead or many trees alive). In such cases an arc sine transformation of data will often prove useful. The arc sine transformation is given by

 $\arcsin(proportion) = \sin^{-1} \sqrt{proportion}$

where proportion is a figure between zero and one

(e.g. the surviving fraction of trees). An important property of the transformation is that the variance near zero or one is stretched out, thus facilitating the analysis of variance (Snedecor & Cochran 1980).

For many growth variables, the variance increases directly with tree size, and the proper transformation is thus the logarithm. In most cases, the natural logarithm (ln) has been applied to achieve a normal distribution of residuals.

Variance inhomogeneity

In the cases where the variance varies with the size of the variable, a transformation of data is the proper way to solve the problem (see above). However, in some cases the provenances simply have different variances irrespective of size, and it is necessary to weight the observations with weights proportional to the reciprocals of the error variances to ensure variance homogeneity (SAS 1988b, cf. Afifi & Clark 1996). There may also be cases where different blocks have different variances, but this has not been observed in the present trial(s).

Weighting occurs in the following sequence: An ordinary analysis of variance of the variable is performed. The residuals from this analysis are grouped according to provenance, and the variance of the residuals for each provenance is calculated. The inverse of these variances is then used as weights in an analysis of variance. When calculating the sums of squares in the model, the weights are multiplied with the squared value of the deviance of each observation from the predicted value (SAS 1988b). This has the effect that provenances with small variances have a larger influence on the model than provenances with larger variances. In other words, the more stable the provenance, the more it counts in the analysis. Provenances with large uncertainty on the other hand have less influence.

4.6 Fixed or random effects

A special problem relates to the choice between considering the effects in the statistical model as fixed or random. Statistically speaking, fixed effects are considered as parameters (unknown constants). Random effects are considered stochastic variables with an expected value of zero and a variance (Skovgård 1994). Fixed effects are used when the individual groups (seedlots) are of interest. Models with random effects are used when interest is in the size of the variation between the groups (described by the variance), including groups that are not represented in the trial. In analysis of random effects it is important that the groups are representative of a larger population of groups, and they should preferably be chosen by randomisation (Skovgård 1994). In the words of Stonecypher (1992), 'fixed models address estimating and testing to infer the existence of true differences among means, whereas the random models address estimating and testing to infer the existence of components of variance'.

To choose between a fixed or a random effects model is a choice with no simple answer. Stonecypher (1992) has formulated the following two questions to facilitate a choice:

- 1. 'Are the conclusion confined to the things actually studied; to the immediate sources of these things; or extended to apply to more general population?'
- 2. 'In complete repetitions of the experiment would the same things be studied again; would new samples be drawn from the same sources; or would new samples be drawn from the general population?'

When the objective is to estimate components of variance, the effects should be considered as random. If the objective is to estimate differences among means, the effects should be considered as fixed. In some cases fixed and random effects may be combined in the model (mixed models). This is the case when special designs are applied, such as split-plot or nested designs.

In our model with only provenances and blocks, it is necessary to choose between considering the provenance effect as random or as fixed. If the aim is to compare the specific provenances and the actual production on the site, it is natural to consider the provenance effect as fixed. If, on the other hand, (i) the provenances are assumed to be representatives of a population of provenances; (ii) the aim is to expand the conclusions to this population; (iii) to estimate the production and (iv) should the experiment be repeated, then the provenance effect should be considered as random.

The results of the statistical tests are irrespective of whether the provenance effect is considered a fixed or a random effect. However, there are major differences in the estimates resulting from the two approaches (see below). Since it may be argued that both the fixed and the random approaches are relevant in this analysis, both sets of estimates have been calculated.

4.7 Test of differences between species and provenances

In our statistical model, differences between provenances for a given trait are tested by an F-test comparing the mean square of provenances with the residual mean square. The hypothesis tested is that there is no difference between the provenances. If the F-test is significant, we reject the hypothesis and conclude that there are significant differences between the provenances.

The testing is done using the GLM procedure in SAS (SAS 1990). Since the testing of random variables may involve combinations of different mean squares (Skovgård 1994), an approximation called Satterthwaites approximation is used in the calculation of degrees of freedom (SAS 1988).

4.8 Lsmeans (estimates from the fixed model)

In the fixed model approach, the estimates for the provenances are calculated as the least square means (lsmeans). The main difference between raw means and lsmeans is that lsmeans account for missing values and imbalanced designs. Thus, in completely balanced designs there are no differences between lsmeans and the raw means. It follows that the lsmeans are the best estimates for the given provenance in the trial.

The confidence intervals and limits are calculated from the formula (Skovgård 1994).

$$X \pm t_{1-lpha/2,(a-1)(b-1)} \sqrt{s^2 / b}$$

where X is the least square mean, α is the confidence level (in this case 0.05, giving a 95% confidence interval), *a* is the number of provenances and b is the number of replicates (blocks) of each provenance. *s*² is the mean square of the error (MS_e). The confidence limits are calculated directly by SAS in the LSMEANS statement with the CL option.

Since the estimates are calculated individually, different provenances may have different lengths of the confidence intervals (due to different variances). In the cases where the data have been weighted, the confidence intervals are adjusted according to the variance of each provenance and thus are of different lengths.

Special problems arise when the data has been transformed. If the least square means and the confidence limits are calculated on basis of the transformed values, the back-transformed estimates will be geometric means rather than arithmetic means. This implies that the estimates become biased towards lower values, and compared to the real values actually are under-estimates. If on the other hand the estimates are calculated using raw data, the lsmeans will be arithmetic means (comparable to the real mean values), but the confidence limits are based on a faulty distribution and will be wrong. In this analysis we have calculated estimates on the transformed values in order to get a fair representation of the differences between provenances. Usually the figures are presented together with a raw mean to circumvent the problem with under-estimation.

4.9 Best Linear Unbiased Predictors (BLUPs - estimates from the random model)

In the random approach, the provenance effects are seen as coming from a normal distribution with an expected value and a variance. This is in opposition to the fixed effect approach, where the provenance effects are seen as constants. Estimating provenance effects in random models is more complicated than in fixed models, because the observed variation between provenances is contemplated as a mixture between true provenance effects and random error variation (*cf.* White & Hodge 1989). The variation between the provenances is therefore always larger than the true 'genetic' variation, except in cases where the error variation is negligible.

In order to predict the effect of a given provenance, it is necessary to correct the estimates for the part of the variance that is due to random error variation. This is done by calculating the best linear unbiased predictors (BLUPs, White & Hodge 1989). The calculation of BLUPs is cumbersome and only feasible with a suitable software package. In this case, the SAS procedure MIXED has been used. It follows from the above that the predicted values for the provenances fall within a smaller range than the least square means. Often the results are presented as deviations from the mean value to allow for easier comparison between different experiments. The deviations are expressed either in real values (m, cm² etc.) or in % deviation from the mean value. Here deviations are presented as % deviations from the mean values.

The problems with transformed values are the same as described for the least square means above. A further complication arises when calculating the deviations from the mean value in percent. If the mean value is calculated on the base of transformed values, and the deviations are calculated on the basis of this back-transformed mean, the deviations from the mean will not sum to zero. In this analysis, we have therefore chosen to base the deviations from the mean value on values calculated after transformation.

The BLUPs are presented with *t*-type confidence intervals. However, these should be interpreted with caution since it is probably wrong to assume that the underlying distribution of the estimates is normal because of the limited sample size (Littell *et al.* 1996). Confidence intervals are presented to give an impression of the variation between the provenances and should not be interpreted with respect to differences between provenances.

5. Results of statistical analysis of individual traits

The below table displays the traits selected for analysis, grouped into growth traits, adaptive traits and quality traits. For a full description of the traits and their calculation, please refer to Annex 4.

Group	Trait description	Analysed trait
Growth	Height growth	Height of tree with diameter corre- sponding to mean basal area (H _G)
	Diameter growth	Diameter of tree corresponding to mean basal area (D _G)
	Mean volume of tree	Average of volumes above bark of trees in plot
	Standing volume per hectare	Volume per hectare
Adaptation	Survival	Survival rate
	Flowering and fruiting	Average score of male flowers
	Foxtailing	Foxtailing percentage
Quality	Stemform	Stemform score (1-9)
	Relative wood density (Pilodyn)	Diameter adjusted pilodyn readings
	Branching	Average branch diameter
	Branching	Average branch/DBH ratio

5.1 Survival

OVERVIEW OF ANALYSIS			
	Aek Nauli	Habinsaran	
Co-variates	None	PLOTY2	
Data transformation required	Yes. Arc sin transformation	Yes. Arc sin transformation	
Weight statement	Yes	Yes	
Outliers	None	None	
F-test	6.27 (***)	3.32 (***)	

Analysis of survival data at year 7 reflects not only differences in 'true' survival rates but is also affected by within-plot competition. For the Aek Nauli trial, irregularities in weeding have resulted in loss of many seedlings, and great care is therefore required in the interpretation of results of this trial.

The Habinsaran trial has a much higher survival rate than the Aek Nauli trial. Many of the seed sources have a survival rate above or close to 90 per cent, e.g. *P. oocarpa* (Mal Paso), Lang Hanh (Vietnam), *P. tecunumanii* (Mt. Pine Ridge), and Bodana (Madagascar). There are not statistical significant differences among the top provenances, but it is possible to distinguish between a high and a low survival group. In the latter one we find *P. patula*, Doi Suthep (Thailand) and Aungban (Myanmar).

The ranking in regard to survival is quite different in the Aek Nauli trial, but as mentioned above, these results have to be interpreted with great care, as also indicated by the very wide confidence intervals. The Lang Hanh (Vietnam), Coto Mines (Philippines) and Doi Suthep (Thailand) have very low survival rates. Especially the low survival rate of Lang Hanh is surprising as it is among the highest ranking provenances in the Habinsaran trial. *P. patula*, *P. tecunumanii* (San Raphael) and Bodana (Madagascar) are high ranked at Aek Nauli, whereas only Bodana is among the best at Habinsaran. *P. patula* and *P. tecunumanii* (San Raphael) have below average survival rates.

Pinus kesiya provenance trial, Aek Nauli, Indonesia

International Series of Pinus kesiya provenance trials, Trial No. 7

Established November 1992. Assessed September 1999

Survival (%)



Pinus kesiya provenance trial, Aek Nauli, Indonesia

International Series of Pinus kesiya provenance trials, Trial No. 7 Established November 1992. Assessed September 1999 Survival. Best linear un-biased predictors (BLUPs)

Expected gain, % deviation from mean

Pinus kesiya provenance trial, Habinsaran, Indonesia

International Series of Pinus kesiya provenance trials, Trial No. 8 Established October 1992. Assessed September 1999

Survival (%)

Pinus kesiya provenance trial, Habinsaran, Indonesia

International Series of Pinus kesiya provenance trials, Trial No. 8 Established October 1992. Assessed September 1999 Survival. Best linear un-biased predictors (BLUPs)

5.2Height growth

OVERVIEW OF ANALYSIS				
	Aek Nauli	Habinsaran		
Co-variates	None	PLOTX and PLOTY2		
Data transformation required	No	No		
Weight statement	No	Yes		
Outliers	Lang Hanh (Block 1)	None		
F-test	12.96 (***)	96.18 (***)		

The analysis of height growth shows highly significant differences among provenances in both trials. The ranking of provenances is very similar in the two trials. The two *P. tecunumanii* sources are at the top in both trials, although the exact ranking of the two sources are different from the one trial to the other.

The *P. tecunumanii* sources are followed by the two sources of *P. oocarpa*. The best source of *P. kesiya* in the Aek Nauli trial is the Coto Mines provenance (Philippines), which is unfortunately not in the Habinsaran trial. Coto Mines is followed by the provenances Doi Suthep (Thailand), Bodana (Madagascar) and Lang Hanh (Vietnam). There are however no statistically significant differences among these sources. At the bottom end we find Nam Now (Thailand), Nong Krating (Thailand) and Aungban (Myanmar). At the very bottom is the Shangsi provenance of *P. yunnanensis*.

In the Habinsaran trial, again the Lang Hanh, Bodana and Doi Suthep are the most promising *P. kesiya* sources. Nam Now, Nong Krating, Aungban and Shangsi sources are at the bottom.

The local *P.merkusid* source is in both trials ranking below the *P. tecunumanii/P. oocarpa* sources but higher than the *P. kesiyd* sources. The *P. patula* source (of Zimbabwe origin) does not show much promise. It is intermediately placed in the Aek Nauli trial, but at the very bottom at Habinsaran. The *P. caribaed* source is only in the Aek Nauli trial, where it is showing a (surprisingly) poor growth.

International Series of Pinus kesiya provenance trials, Trial No. 7 Established November 1992. Assessed September 1999

Pinus kesiya provenance trial, Aek Nauli, Indonesia

International Series of Pinus kesiya provenance trials, Trial No. 7 Established November 1992. Assessed September 1999 Height gain. Best linear un-biased predictors (BLUPs)

Expected gain, % deviation from mean

Pinus kesiya provenance trial, Habinsaran, Indonesia

International Series of Pinus kesiya provenance trials, Trial No. 8

Pinus kesiya provenance trial, Habinsaran, Indonesia

International Series of Pinus kesiya provenance trials, Trial No. 8 Established October 1992. Assessed September 1999 Height gain. Best linear un-biased predictors (BLUPs)

5.3 Diameter growth

	OVERVIEW OF ANALYSIS	S
	Aek Nauli	Habinsaran
Co-variates	None	PLOTX, PLOTY and PLOTY2
Data transformation required	No	No
Weight statement	No	Yes
Outliers	Lang Hanh (Block 1)	None
F-test	6.29 (***)	28.31 (***)

There are highly significant differences in regard to diameter growth in both trials.

The results of the Aek Nauli trial- again - have to be interpreted with care as differences in survival between plots may have influenced diameter growth. This is reflected in the considerably larger confidence intervals in the Aek Nauli trial compared to the Habinsaran trial.

If we look at the results of the Habinsaran trial first, the ranking of provenances is not very different from what we have seen for height growth. We have the *P. tecunumanii* sources at the top, followed by *P. oocarpa*. Lang Hanh (Vietnam) and Bodana (Madagascar) are again best among the *P. kesiya* sources, although there are only small and statistically insignificant differences among the best *P. kesiya* sources. The *P. yunnanensis* source (Shangsi-China), *P. patula* and Aungban (Myanmar) are at the bottom end and they can be distinguished also statistically from the above mentioned sources.

In the Aek Nauli trial the ranking is quite different, which is believed to a large extent due to the differences in survival. Doi Suthep (Thailand) and Lang Hanh (Vietnam) placed in the top together with the two *P. tecunumanii* sources. The two *P. oocarpa* sources are ranked relatively low. In the lower end, the picture is identical to the Habinsaran trial with Shangsi (China), Aungban (Myanmar) and *P. patula* having the poorest diameter growth.

Pinus kesiya provenance trial, Aek Nauli, Indonesia

International Series of Pinus kesiya provenance trials, Trial No. 7 Established November 1992. Assessed September 1999

Pinus kesiya provenance trial, Aek Nauli, Indonesia

International Series of Pinus kesiya provenance trials, Trial No. 7 Established November 1992. Assessed September 1999 Diameter gain. Best linear un-biased predictors (BLUPs)

Expected gain, % deviation from mean

Pinus kesiya provenance trial, Habinsaran, Indonesia

International Series of Pinus kesiya provenance trials, Trial No. 8 Established October 1992. Assessed September 1999

Diameter (cm)

Pinus kesiya provenance trial, Habinsaran, Indonesia

International Series of Pinus kesiya provenance trials, Trial No. 8 Established October 1992. Assessed September 1999 Diameter gain. Best linear un-biased predictors (BLUPs)

5.4 Mean volume of tree

	OVERVIEW OF AN	ALYSIS
	Aek Nauli	Habinsaran
Co-variates	None	PLOTX and PLOTY
Data transformation required	No	No
Weight statement	No	Yes
Outliers	None	None
F-test	6.63 (***)	52.14 (***)

Mean volume of tree is calculated as the average of the volumes of individual trees. As both height and diameter are included in the volume formula, the trait thus illustrates a combined effect of height and diameter.

Again, the Aek Nauli results have to be interpreted with care because of the different survival rates and consequently diameter growth.

In the Habinsaran trial, the two *P. tecunumanii* seed sources are at the top, followed by the *P. oocarpa* sources. They again are followed by the local *P. merkusii* and only then we find the best *P. kesiya* sources. They are Lang Hanh (Vietnam) and Bodana (Madagascar). *P. patula* and *P. yunnanensis* are at the bottom.

In the Aek Nauli trial, we also have the two *P. tecunumanii* sources at the top, but they are followed closely by Doi Suthep (Thailand) and Coto Mines (Philippines). At the bottom we find, as in the Habinsaran trial, Aungban (Myanmar) and Shangsi (China).

Pinus kesiya provenance trial, Aek Nauli, Indonesia

International Series of Pinus kesiya provenance trials, Trial No. 7

Established November 1992. Assessed September 1999

Volume of mean tree (m3)

Pinus kesiya provenance trial, Aek Nauli, Indonesia

International Series of Pinus kesiya provenance trials, Trial No. 7 Established November 1992. Assessed September 1999

Expected gain, % deviation from mean

Pinus kesiya provenance trial, Habinsaran, Indonesia

International Series of Pinus kesiya provenance trials, Trial No. 8 Established October 1992. Assessed September 1999

accondition October 1992, Assessed September 15

Pinus kesiya provenance trial, Habinsaran, Indonesia

International Series of Pinus kesiya provenance trials, Trial No. 8 Established October 1992. Assessed September 1999 Volume of mean tree. Best linear un-biased predictors (BLUPs)

	OVERVIEW OF ANALYSIS	
	Aek Nauli	Habinsaran
Co-variates	PLOTXY	PLOTX and PLOTY2
Data transformation required	No	No
Weight statement	Yes	Yes
Outliers	None	None
F-test	6.01 (***)	33.88 (***)

5.5 Total volume per hectare

The analysis of total volume production can be seen as an analysis summarising survival, height growth and diameter growth in one analysis as all three traits are included in the calculation.

The Habinsaran trial shows a ranking of provenances almost identical to what we have seen for mean volume of tree. The two *P. tecunumanii* sources are by far the most productive, with the San Raphael source slightly better than the Mt. Pine Ridge although the difference is not statistically significant. Following the *P. tecunumanii* sources, but with significantly lower production, we have the two *P. oocarpa* provenances followed by *P. merkusii*. The best sources of *P. kesiya* are again Bodana (Madagascar) and Lang Hanh (Vietnam).

In the Aek Nauli trial, we also have the *P. tecunu-manii* sources at top, but the ranking underneath is somewhat different to the Habinsaran trial. The Bodana landrace from Madagascar is the best *P. kesiyd* source like in the Habinsaran trial, whereas

International Series of Pinus kesiya provenance trials, Trial No. 7

Established November 1992. Assessed September 1999

Volume per ha (m3/ha)

Pinus kesiya provenance trial, Aek Nauli, Indonesia

International Series of Pinus kesiya provenance trials, Trial No. 7 Established November 1992. Assessed September 1999 Volume per hectare. Best linear un-biased predictors (BLUPs)

Expected gain, % deviation from mean

Pinus kesiya provenance trial, Habinsaran, Indonesia

International Series of Pinus kesiya provenance trials, Trial No. 8 Established October 1992. Assessed September 1999

Volume per hectare (m3/ha)

Pinus kesiya provenance trial, Habinsaran, Indonesia

International Series of Pinus kesiya provenance trials, Trial No. 8 Established October 1992. Assessed September 1999 Volume gain. Best linear un-biased predictors (BLUPs)

5.6 Stemform

OVERVIEW OF ANALYSIS			
	Aek Nauli	Habinsaran	
Co-variates	PLOTY2	None	
Data transformation required	Yes. Ln transformation	No	
Weight statement	Yes	Yes	
Outliers	Nong Krating (Block 2)	None	
F-test	8.05 (***)	11.60 (***)	

The statistical analysis reveals significant differences among provenances in regard to stemform in both trials.

The *P. merkusii* and the *P. caribaea* source (the latter is only in the Aek Nauli trial) have a considerably poorer stemform than the other sources.

The ranking of provenances is different in the two trials. In the Habinsaran trial, *P. patula* is at the top followed by *P. oocarpa* (Honduras) and the two *P. tecunumanii* sources. The best *P. kesiya* sources are Nong Krating (Thailand), Nam Now (Thailand) and Bodana (Madagascar). Aungban (Myanmar), Lang Hanh (Vietnam), Shangsi (China) and Doi Suthep (Thailand) have the poorest stemform among the *P. kesiya* sources.

At Aek Nauli, Nam Now (Thailand) and Doi Suthep (Thailand) are at the top. This in contrast to the Habinsaran trial where Nam Now is intermediate, and Doi Suthep is in the lower half. Next at Aek Nauli we have the sources of *P. oocarpa*, *P. tecunumanii*, *P. patula* and Bodana (Madagascar) which were at the very top at Habinsaran.

For both trials, there are only small – and not statistically significant – differences among the top ranking seed sources.

Pinus kesiya provenance trial, Aek Nauli, Indonesia

International Series of Pinus kesiya provenance trials, Trial No. 7 Established November 1992. Assessed September 1999

Pinus kesiya provenance trial, Habinsaran, Indonesia

International Series of Pinus kesiya provenance trials, Trial No. 8 Established October 1992. Assessed September 1999 Stemform (1-9 score)

Pinus kesiya provenance trial, Aek Nauli, Indonesia

International Series of Pinus kesiya provenance trials, Trial No. 7 Established November 1992. Assessed September 1999 Stemform. Best linear un-biased predictors (BLUPs)

Pinus kesiya provenance trial, Habinsaran, Indonesia

International Series of Pinus kesiya provenance trials, Trial No. 8 Established October 1992. Assessed September 1999 Stemform. Best linear un-biased predictors (BLUPs)

5.7 Wood density (Pilodyn)

OVERVIEW OF ANALYSIS			
	Aek Nauli	Habinsaran	
Co-variates	None	PLOTY2	
Data transformation required	Yes. Square-root transformation	Yes. Ln transformation	
Weight statement	No	Yes	
Outliers	None	None	
F-test	3.71(***)	7.68 (***)	

Diameter adjusted pilodyn readings (ref. Annex 4) are used in the analysis.

Both the Aek Nauli and the Habinsaran trial reveal significant differences between provenances. The ranking of provenances, however, is different in the two trials. As we have seen for most other traits, the confidence intervals in the Aek Nauli trial are much wider than in the Habinsaran trial.

The sources with the fastest growth (diameter and total volume production), i.e. the *P. tecunumanii* and *P. oocarpa* sources, have the highest pilodyn values corresponding to the lowest wood densities. Consequently, the slow growing sources have in general the highest wood densities. This is the general picture, but there are exceptions, and the exact ranking, as mentioned above, is different in the two trials.

Most remarkable is *P. patula* that is ranked at the top at Habinsaran, but at the bottom end in the Aek Nauli trial. A ranking among the top sources is what would be expected considering the poor growth of *P. patula* in both trials. Also the Aungban (Myanmar) and *P. oocarpa* (Mal Paso) sources have very different rankings in the two trials.

There are significant differences when comparing the top and bottom, but differences among many of the sources ranked in-between are only small, and not statistically significant.

Pinus kesiya provenance trial, Aek Nauli, Indonesia International Series of Pinus kesiya provenance trials, Trial No. 7

Pinus kesiya provenance trial, Habinsaran, Indonesia

International Series of Pinus kesiya provenance trials, Trial No. 8

Pinus kesiya provenance trial, Aek Nauli, Indonesia

International Series of Pinus kesiya provenance trials, Trial No. 7 Established November 1992. Assessed September 1999 Pilodyn, Best linear un-biased predictors (BLUPs)

Pinus kesiya provenance trial, Habinsaran, Indonesia

International Series of Pinus kesiya provenance trials, Trial No. 8 Established October 1992. Assessed September 1999 Pilodyn. Best linear un-biased predictors (BLUPs)

5.8 Branching (branch diameter)

	OVERVIEW OF ANALYSIS	
	Aek Nauli	Habinsaran
Co-variates	PLOTX2	PLOTY
Data transformation required	No	No
Weight statement	Yes	Yes
Outliers	None	None
F-test	11.87(***)	4.27 (***)

Branch diameter (largest branch in whorl at 1/10 of tree height) has only been assessed on the *P. kesiyd* /*P. yunnanensis* sources, and therefore the analysis is restricted to these sources. The analysis reveals significant differences between provenances in both trials.

In both trials, the Shangsi (*P. yunnanensis*) source has considerably smaller branch diameters than the other sources. The growth potential of this source is very poor, and the small branches are related to this fact.

The results of the Aek Nauli trial, again, have to be interpreted with care as low survival most likely has influenced development of thick branches. This is e.g. the case for Doi Suthep (Thailand) which in the Habinsaran trial is among the provenances with smallest branch diameter but at the very top at Aek Nauli.

If we leave out Shangsi, there are not great differences among the sources, and differences are not statistically significant. There is maybe a slight tendency to provenances having better growth also having thicker branches.

Pinus kesiya provenance trial, Aek Nauli, Indonesia

International Series of Pinus kesiya provenance trials, Trial No. 7

Pinus kesiya provenance trial, Aek Nauli, Indonesia

International Series of Pinus kesiya provenance trials, Trial No. 7 Established November 1992. Assessed September 1999 Branch diameter. Best linear un-biased predictors (BLUPs)

Pinus kesiya provenance trial, Habinsaran, Indonesia International Series of Pinus kesiya provenance trials, Trial No. 8 Established October 1992. Assessed September 1999

Pinus kesiya provenance trial, Habinsaran, Indonesia

International Series of Pinus kesiya provenance trials, Trial No. 8 Established October 1992. Assessed September 1999 Branch diameter. Best linear un-biased predictors (BLUPs)

5.9 Foxtailing

OVERVIEW OF ANALYSIS								
	Aek Nauli	Habinsaran						
Co-variates	None	None						
Data transformation required	No	No						
Weight statement	Yes	Yes						
Outliers	Block 1: P. tecunumanii (MPR), Lang Hanh and Doi Suthep; Block 2: Nong Krating; Block 3: Doi Suthep and Shangsi; and Block 4: Lang Hanh	None						
F-test	1.45(NS)	8.91 (***)						

Frequency of foxtails has only been assessed on the *P. kesiya/P. yunnanensis* sources.

There are not significant differences among sources in the Aek Nauli trial, whereas in the Habinsaran trial there are significant differences.

It looks as *P. yunnanensis* (Shangsi) has significantly fewer foxtails than the *P. kesiya* sources. For the *P. kesiya* sources the frequency of foxtails is high; between 55 and 75 percent. There are however not statistically significant differences among provenances. There is a tendency, although not statistically significant, to Thai sources having fewer foxtails than the other sources.

Foxtailing is an important trait to consider. First of all it gives an indication of the adaptability of the source to the site (a low frequency is in general an indication of good adaptation). Moreover, and probably of greater practical importance, foxtails will influence quality. Foxtails will result in ramicorns, i.e. thick branches growing (co-evolving) vertically along with the main stem as there are no branches on the foxtail to suppress this growth. Thick branches mean lower quality, especially if timber is the final products. In addition, harvesting operations become more difficult. Finally, foxtails will often result in broken stems because of the soft wood.

Pinus kesiya provenance trial, Aek Nauli, Indonesia

International Series of Pinus kesiya provenance trials, Trial No. 7

Established November 1992. Assessed September 1999

Pinus kesiya provenance trial, Habinsaran, Indonesia

International Series of Pinus kesiya provenance trials, Trial No. 8 Established October 1992. Assessed September 1999

Foxtail percentage (%)

Pinus kesiya provenance trial, Aek Nauli, Indonesia

International Series of Pinus kesiya provenance trials, Trial No. 7 Established November 1992. Assessed September 1999 Foxtailing. Best linear un-biased predictors (BLUPs)

Pinus kesiya provenance trial, Habinsaran, Indonesia

International Series of Pinus kesiya provenance trials, Trial No. 8 Established October 1992. Assessed September 1999 Foxtailing. Best linear un-biased predictors (BLUPs)

5.10 Flowering

OVERVIEW OF ANALYSIS									
Aek Nauli Habinsaran									
Co-variates	LEVEL, PLOTX and PLOTY	None							
Data transformation required	No	No							
Weight statement	Yes	Yes							
Outliers	Block 1: P. tecunumanii (MPR), Lang Hanh and Doi Suthep; Block 2: Nong Krating; Block 3: Doi Suthep and Shangsi; and Block 4: Lang Hanh	None							
F-test	3.52(***)	0.70 (NS)							

Assessment of flowering has only been done on the *Pinus kesiya*/*P. yunnanensis* sources.

Frequency of flowers provides an indication of the adaptation of the sources to site. A good flowering and fruiting will generally be interpreted as a sign of good adaptation to the site, and vice versa.

The two trials are quite young, only about 7 years of age, and flowering and fruiting may have only just commenced. Consequently, male flowers were the only development stage that was present on most trees, and hence the only stage that could be assessed and analysed.

In the Habinsaran trial, there is very sparse flowering, and no statistical differences between the provenances.

The Aek Nauli trial has more frequent male flowering and there are significant provenance differences. *P. yunnanensis* (Shangsi) and Aungban (Myanmar) have the lowest flowering scores, and Doi Suthep (Thailand) the highest. It is a question if not the uneven survival rates in the Aek Nauli trial have an influence here. The uneven survival rates may have lead to different light conditions in plots which have facilitated flowering to a variable degree. The differences may thus more be because of survival differences than 'true' provenance differences.

Pinus kesiya provenance trial, Aek Nauli, Indonesia

Pinus kesiya provenance trial, Aek Nauli, Indonesia International Series of Pinus kesiya provenance trials, Trial No. 7

Established November 1992. Assessed September 1999 Male flowers. Best linear un-biased predictors (BLUPs)

Pinus kesiya provenance trial, Habinsaran, Indonesia International Series of Pinus kesiya provenance trials, Trial No. 8 Established October 1992. Assessed September 1999 Male flowers (1-9 score)

No figure of BLUP-estimates for the Habinsaran trial as there are no statistical differences between provenances.

6. Conclusions

Conclusions are to be based mainly on the performance in the Habinsaran trial. This is because of the many lost seedlings in the Aek Nauli trial that has influenced the growth and development. Differences in ranking of seedlots in the two trials are believed to be mainly an effect of this, and not differences caused by different environments at the two sites (i.e. not believed to be genotype x environment interaction).

6.1 Growth

Growth traits should be given key importance in the interpretation of trial results, as production of pulpwood has high priority.

The analysis shows not much promise for the tested sources of P. kesiya in comparison to the included controls - most of the tested controls have a considerably faster growth than the P. kesiya sources.

The two *P. tecunumanii* sources are superior to the other sources in regard to growth in both trials, and is the most promising species at sites similar to the test sites. The San Raphael provenance has a slightly better growth than the Mt. Pine Ridge provenance, but differences are not statistically significant. It would be interesting to test more sources of *P. tecunumanii*, as other sources may have an even greater potential. The company has such trials under way. Mean annual production – based on the results of the Habinsaran trial – is approx. 40 m3/ha/year.

Following *P. tecunumanii* we find the two sources of *P. oocarpa*, but growth is considerably slower than that of *P. tecunumanii*. In the Aek Nauli trial the two *P. oocarpa* sources have almost the same growth, whereas at Habinsaran, the Honduran source has the fastest growth.

Next in ranking is the local (Indonesian) *P. merkusii* source, and only then we arrive at the best *P. kesiya* sources.

Lang Hanh (Vietnam) and Bodana (Madagascar) are the best growth performers among the *P. kesiya* sources. It is interesting to note that the Madagascar source (landrace) almost certainly originates from the Central plateau of Vietnam (Armitage & Burley, 1980), i.e. from the same region as the Lang Hanh seedsource. It was introduced from here to Madagascar in the 1920's.

Coto Mines (Philippines) is performing well in the Aek Nauli trial (height growth) but is unfortunately not included at Habinsaran.

Doi Suthep (Thailand) may be mentioned together with the above sources, mainly based on a relatively good growth in the Aek Nauli trial. It seems, however, that this source has a low survival rate. *P. patula* shows little promise based on the two trials. It is very slow growing, has a low survival rate, and generally looks unhealthy.

The poorest growth performer is the *P. yunnanensis* source that has a volume production less than one fifth of the *P. tecunumanii* sources.

6.2 Adaptation

Adaptive traits include survival percentage, foxtail frequency and flowering. The two latter traits have only been assessed on the *P. kesiya/P. yunnanensis* sources.

There are no statistical significant differences among top ranking sources in regard to survival. The best *P. kesiya* performers in regard to growth, Lang Hanh (Vietnam) and Bodana (Madagascar), also have a good survival. In the other end of the scale, *P. patula*, Doi Suthep (Thailand) and Aungban (Myanmar) have a low survival rate.

The *P. yunnanensis* source has a considerably lower frequency of foxtails than the *P. kesiya* sources, but the result has little practical value because of the extremely poor growth of this source. The high frequency of foxtails for the *P. kesiya* sources may be another constraint for a more intensive use of the species.

6.3 Quality

Quality parameters are stemform, wood density (pilodyn) and branch diameter. The *P. tecunumaniil* sources, the *P. oocarpal* ones and *P. patulal* have the best stemform. *P. merkusiil* has a considerably poorer stemform than the rest of sources, with the *P. kesiyal* sources forming an intermediate group. The Bodana (Madagascar) source is again among the best, whereas the Lang Hanh (Vietnam) has a poorer stemform.

The more slow growing sources generally have a better wood density (esp. Shangsi and *P. merkusii*) and thinner branches than the faster growing sources (esp. *P. tecunumanii*) and *P. oocarpa*). The Lang Hanh and Bodana sources are again here among the highest ranking sources of *P. kesiya*.

6.4 Conclusion

Based on the performance in the two trials, the two sources of *P. tecunumanii* are the most promising. Further testing and investigation of the genetic variation within this species is recommended. The Research & Development Division has trials under way with additional sources of *P. tecunumanii* and these trials will provide valuable information on the most appropriate sources of the species.

P. kesiya shows little promise as a plantation species on the tested sites. The growth is much slower than P. tecunumanii and P. oocarpa, and it also compares less favorable in regard to stemform. P. kesiya may have a larger potential on poorer and harsher sites more influenced by fires (Clegg, pers. comm.). If results of the present trials also are applicable under such conditions, the analysis indicates the Lang Hanh source (natural population from the central plateau of Vietnam) and Bodana A8 (offspring from seed orchard in Madagascar, material probably originally from Vietnam) as the most promising sources. The Lang Hanh source may show an even larger potential in subsequent generations as inbreeding depression from the natural population breaks down. Other sources from the Central plateau of Vietnam could also be of potential interest. The same holds for Philippine sources, which are only represented with one provenance and only in the Aek Nauli trial.

7. References

Afifi, A.A. and V. Clark. 1996. Computer-aided multivariate analysis. Chapman & Hall, London (3rd Edition).

- Armitage, F.B. and J. Burley. 1980. *Pinus kesiya*. Tropical Forestry Papers No. 9. Department of Forestry, Commonwealth Forestry Institute, University of Oxford, Oxford, U.K. 199p
- Barnes, R.D., H. Keiding and G.L. Gibson. 1989. Progress on seed collection for the second stage international provenance trials of *Pinus kesiya*. Forest Genetic Resources Information No. 17.
- Box, G.E.P., W.G. Hunter and J.S. Hunter. 1978. Statistics for experimenters. John Wiley and Sons Inc., New York.
- Burley, J. and P.J. Wood. 1976. Manual on Species and Provenance Research with particular reference to the tropics. Tropical Forestry Papers No. 10. Department of Forestry, Commonwealth Forestry Institute, University of Oxford, Oxford, U.K. 226 p.
- DFSC 1996. Proposal for a common evaluation of an international series of *Pinus kesiya* trials. DFSC Working Paper. 8 p.
- DFSC 1997. Status on *Pinus kesiya* international provenance trials Proposed Work Program 1996-99. DFSC Working Paper. Danida Forest Seed Centre, Humlebæk, Denmark. 10 p.
- DFSC 1998. International Series of Provenance Trials of *Pinus kesiya*. Field Assessment manual. Danida Forest Seed Centre, Humlebaek, Denmark. 19 p.
- FAO 1988. Report of the Sixth Session of the FAO Panel of Experts on Forest Gene Resources. Food and Agriculture Organization of the United Nations. Rome, Italy. 79 p.
- Gibson, G.L. and R.D. Barnes. 1984. Status of the international provenance trials of *Pinus kesiyal* and alternatives for future development. In: Barnes, R.D. and Gibson, G.L. (Eds.): Provenance and genetic improvement strategies in tropical forest trees. Proceedings of the IUFRO Conference, Mutare, Zimbabwe, April 1984. Commonwealth Forest Institute and Forest Research Centre, Harare.
- Keiding, H., H. Wellendorf and E.B. Lauridsen. 1986. Evaluation of an international series of teak provenance trials. Danida Forest Seed Centre, Humlebaek, Denmark. 81 p.
- Littell, R.C., G.A. Milliken, W.W. Stroup, R.D. Wolfinger. 1996. SAS® System for mixed models. SAS Institute Inc., Cary, NC, 633 pp.
- SAS 1988a. SAS® Procedures Guide, Release 6.03 Edition. SAS Institute Inc., Cary, NC, 441 pp.
- SAS 1988b. SAS/STAT® Users Guide, Release 6.03 Edition. SAS Institute Inc., Cary, NC, 1928 pp.
- SAS 1991. SAS® System for Statistical Graphics, First Edition. SAS Institute Inc., Cary, NC, 697 pp.
- Skovgård, I.M. 1994. Statistisk Forsøgsplanlægning (in Danish). Jordbrugsforlaget, Copenhagen, Denmark, 318 pp.

Skovgaard, I.M. and P. Brockhoff. 1998. Multivariate analysis and variance components. Lecture notes, Dept. of Mathematics and Physics, The Royal Veterinary and Agricultural University, Copenhagen, 41 pp.

Snedecor, G.W. and W.G. Cochran. 1980. Statistical methods. Iowa State University Press, 7th ed., 507 pp. Stonecypher, R.W. 1992. Computational methods. *In*: Fins, L., Friedman, S.T. & Brotschol, J.V. (eds.):

- Handbook of quantitative forest genetics. Kluwer Academic Publishers, Dordrecht, pp. 195-228. White, T.L. and G.R. Hodge. 1989. Predicting breeding values with applications in forest tree improve-
- ment. Kluwer Academic Publishers, Dordrecht, 367 pp.

Personal communication

Brockhoff, P. Lecturer in statistics. The Royal Veterinary and Agricultural University, Copenhagen.

Clegg, P.A. Research manager, RGMI Forestry, Research & Development Division.

Annexes

Annex 1. Maps

Aek Nauli

Local ID	DFSC Acc. No.	Species	Provenance	Country
712	-	P. oocarpa	Mal Paso	Guatemala
713	-	P. tecunumanii	Mt. Pine Ridge	Belize
714	-	P. oocarpa	El Paraiso	Honduras
715	-	P. tecunumanii	San Raphael	Nicaragua
716	-	P. caribaea	Guanaja	Honduras
718	1572/85	P. kesiya	Coto Mines	Philippines
719	1525/85	P. kesiya	Nam Now	Thailand
720	1521/85	P. kesiya	Nong Krating	Thailand
721	1519/85	P. kesiya	Lang Hanh	Vietnam
722	1522/85	P. kesiya	Doi Suthep	Thailand
723	1523/85	P. kesiya	Doi Inthanon	Thailand
724	1639/86	P. kesiya	Simao	China
725	1783/88	P. kesiya	Bodana A8	Madagascar
726	1773/88	P. kesiya	Aungban	Myanmar
727	1633/86	P. yunnanensis	Shangsi	China
183	-	P. merkusii	Indonesia	Indonesia
118	-	P. patula	Zimbabwe landrace	Zimbabwe
366	-	E. grandis	Coff's Harbour	Australia

Habinsaran

1				l	1 5	5	6	7	8	q			
1	\		14	5 7	12 7	'13 [']	714	715	719	720			
	\mathbf{i}		14	7	26 7	25	183	723	722	721			
		N	13	3 7	27			I					
			12	2 1	18								
	~		11	3	66	_							
	- -		10	726	722						X X 4.1 4.2	X 4.3	X 4.4
	io Dro	п	9	723	713						X X	X 33	X 3.4
			8	712	727						X X 21 22	X 2.3	X 2 4
			7	715	366						X X	X	X
6	719	721	712	720	725								
5	725	183	714	719	721]							
4 III	726	720	722	714	118	183	30	56			Planted	=	Oct ' 1992
3	118	366	715	724	183	118	71	3	IV		Spacing	=	3 x 2 m
2	727	713	723	723	719	721	72	22			Large	=	1.0 Ha
1	727	714	715	725	726	720	71	2					
	I	2	3_	4	5	6	F	,					
			_ PL	.0TX									

Local ID	DFSC Acc. No.	Species	Provenance	Country
712	-	P. oocarpa	Mal Paso	Guatemala
713	-	P. tecunumanii	Mt. Pine Ridge	Belize
714	-	P. oocarpa	El Paraiso	Honduras
715	-	P. tecunumanii	San Raphael	Nicaragua
716	-	P. caribaea	Guanaja	Honduras
718	1572/85	P. kesiya	Coto Mines	Philippines
719	1525/85	P. kesiya	Nam Now	Thailand
720	1521/85	P. kesiya	Nong Krating	Thailand
721	1519/85	P. kesiya	Lang Hanh	Vietnam
722	1522/85	P. kesiya	Doi Suthep	Thailand
723	1523/85	P. kesiya	Doi Inthanon	Thailand
724	1639/86	P. kesiya	Simao	China
725	1783/88	P. kesiya	Bodana A8	Madagascar
726	1773/88	P. kesiya	Aungban	Myanmar
727	1633/86	P. yunnanensis	Shangsi	China
183	-	P. merkusii	Indonesia	Indonesia
118	-	P. patula	Zimbabwe landrace	Zimbabwe
366	-	E. grandis	Coff's Harbour	Australia

Annex 2. Trial descriptions Aek Nauli, Indonesia

TRIAL ESTABLISHMENT AND MANAGEMENT

Year and month of establishment: November 1992

Area (ha): 1.0 ha

Initial spacing (m x m): 3 m x 3 m

Soil preparation (time, method/intensity): Site ripper mounded

Planting method(age of seedlings, type): Polybags, 6 cm diameter, 10 cm height, seedlings probably 25-30 cm

Beating up (time, %): No information

Irrigation (time, amount): None

Fertilization (time, type, amount): No information

Weeding (time, intensity): Irregular, none in 1993-94

Thinning (time, intensity): None

Firelines: None

TRIAL DESIGN

Statistical design: Randomized complete block design

No. of replications (blocks): 4 replications

No. of treatments (provenances): 16 provenances (see list in Annex 1)

Plot size (No. of trees in plot): $16 (4 \times 4)$

Demarcation (blocks, plots): Labels, poles in plot corners. Note problems with identification of seedlot 723 (Doi Inthanon, Thailand)(not kesiya).

PROTECTION STATUS

Status (describe any disturbances/damages): Survival is generally low. Many plots with no surviving trees. Many trees are believed to have been cut by accident when undertaking weeding (irregular weeding). Maybe also by pesticide application...

Guarding (permanent, regular, none): Permanent guarding. Trial is close to office and R&D nursery

Habinsaran, Indonesia

TRIAL ESTABLISHMENT AND MANAGEMENT

Year and month of establishment: October 1992

Area (ha): 1.0 ha

Initial spacing (m x m): 3 m x 3 m

Soil preparation (time, method/intensity): Site ripper mounded

Planting method(age of seedlings, type): Polybags, 6 cm diameter, 10 cm height, seedlings probably 25-30 cm

Beating up (time, %): No information

Irrigation (time, amount): None

Fertilization (time, type, amount): No information

Weeding (time, intensity): No information

Thinning (time, intensity): None

Firelines: None

TRIAL DESIGN

Statistical design: Randomized complete block design

No. of replications (blocks): 4 replications

No. of treatments (provenances): 15 provenances (see list in Annex 1)

Plot size (No. of trees in plot): 16 (4 x 4)

Demarcation (blocks, plots): Labels, poles in plot corners. Note problems with identification of seedlot 723 (Doi Inthanon, Thailand)(not kesiya).

PROTECTION STATUS

Status (describe any disturbances/damages): Good survival and growth. Problems with proper identification in some plots. Seedlot 723, Doi Inthanon is not P. kesiya but an un-identified source of P. tecunumanii. Problems in other plots identified as well....

Guarding (permanent, regular, none): Regular guarding. Trial is 2.5 km from sector office

Annex 3.

Site description – Aek Nauli

LOCATION

Province: North Sumatra District: Simalungun Latitude (degrees and minutes): 02°44'04''N Longitude (degrees and minutes): 98°53'39E Altitude (m above sea level): 1250 m above sea level

Managing office/institution: PT INTI Indorayon Utama, R&D Department Owner: do User(s): do

Distance to nearest office responsible for management of the trial (km): 70 Distance to nearest villages/towns (km): *Ujung Mauli, 2 km* Number of inhabitants in the nearest villages/towns: *approx. 200*

Type of area (e.g. research station, managed forest, etc.): *Managed forest plantations, mainly Eucalyptus sp. for pulp*

CLIMATE

Nearest weather station:

Name of the station: *Aek Nauli base camp (9 km E of trial site)* Latitude (degrees and minutes): *see above* Longitude (degrees and minutes): *see above* Altitude (m a.s.l.): *1200 m above sea level*

Climatic data ¹	Jan.	Feb.	Mar.	Apr.	May	June	July	Aug.	Sep.	Oct.	Nov.	Dec.	Year
Rainfall (mm)	147.5	152.9	174.5	205.9	173.9	143.7	138.9	191.7	173.4	177.6	214.1	220.0	2114.2
Temp. mean (°C)	20.5	20.8	20.9	21.3	21.9	21.9	21.5	21.4	21.1	20.7	20.8	22.8	21.1
Temp. mean max. ² (°C)	-	-	-	-	-	-	-	-	-	-	-	-	-
Temp. mean min. ³ (°C)	-	-	-	-	-	-	-	-	-	-	-	-	-
Evapotranspiration ⁴ (mm)	-	-	-	-	-	-	-	-	-	-	-	-	-

¹ Period of observations 1988-98 ² Average of daily maximum temperatures

³ Average of daily minimum temperatures ⁴ Potential evapotranspiration (ETP) - Penman's formula

Rainy season:

Number/type of seasons:		one		two	x	Even
-------------------------	--	-----	--	-----	---	------

Period(s):	(specify months)

Length of rainy season:

No. of intermediate days:(pre- and posthumid period of the growing season)No. of wet days:(growing season)

Number of dry months per year (< 50 mm rain/month): *None* Frost (number of days/year): *None* Prevailing winds (direction, period, speed): *W to SW very occasionally strong* Alternative weather station: Name of the station: Latitude (degrees and minutes): Longitude (degrees and minutes): Altitude (m a.s.l.):

Climatic data ¹	Jan.	Feb.	Mar.	Apr.	May	June	July	Aug.	Sep.	Oct.	Nov.	Dec.	Year
Rainfall (mm)													
Temp. mean (°C)													
Temp. mean max. ² (°C)													
Temp. mean min. ³ (°C)													
Evapotranspiration ⁴ (mm)													
¹ Period of observations ³ Average of daily minimum temperatures ⁴ Potential evapotranspiration (ETP) - Penman's formula Rainy season:													
Number/type of seasons: one two even/irregular													
	Pe	riod(s)	:	(specif	fy month	s)							
Len	gth of 1 No No	rainy se o. of in o. of w	eason: itermedi et days:	ate day	vs: (growi	(pre ng seas	- and p son)	oosthui	nid per	riod of	the gro	owing s	eason)
Number of dry months per year (< 50 mm rain/month): Frost (number of days/year): Prevailing winds (direction, period, speed):													
TOPOGRAPHY (slope) of trial site													
Flat/gentle (0-8%) x Intermediate (9-30%) Steep (>30%)													

GENERAL SOIL DESCRIPTION

Soil texture		Soil depth	Soil drainage/ n		Gravel content, topsoil		
1. Light/sandy		1. Shallow (< 50 cm)	Х	1. Well drained	Х	1. None (< 15 %)	
2. Medium/loamy		2. Deep (50-100 cm)		2. Seasonal		2. Gravelly (15-35 %)	Х
3. Heavy/clayey	X	3. Very deep (>100 cm)		3. Permanent		3. Stony (> 35 %)	

Organic matter content	Reaction (pH)	Soil salinity		Groundwater			
1. Poor (< 2 % DM)	Х	1. Acid (pH < 6.5)	X	1. None	Х	1. Shallow (< 50 cm)	
2. Medium (2-5 % DM)		2. Neutral (6.5-7.5)		2. Moderate		2. Deep (50 - 150 cm)	
3. Rich (> 5 %)		3. Alkaline (pH> 7.5)		3. High		3. Very deep (>150 cm)	Х

Specify soil unit, soil association and phases (subdivisions of soil units) according to the Soil map of the world (FAO-Unesco 1971-1979), if known:

Natural (original) vegetation type: Natural forest

Dominant natural (original) genera/species: Many

Land use history: Natural forest -> degraded forest/scrub -> pines (+/- 50 years) -> eucalypts (4 yrs)

RESULTS OF SOIL SAMPLE

Results of the laboratory analysis of the soil samples taken at the trial site.

The variables are:

Soil sample depth
Particle size less than 2 µm (0.002 mm) in diameter
Particle size between 2 and 20 µm (0.002 - 0.020 mm) in diameter
Particle size between 20 and 63 μ m (0.020 - 0.063 mm) in diameter
Particle size between 63 and 125 μ m (0.063 - 0.125 mm) in diameter
Particle size between 125 and 250 µm (0.125 - 0.250 mm) in diameter
Particle size between 250 and 500 μ m (0.250 - 0.500 mm) in diameter
Particle size between 500 and 2000 μ m (0.500 - 2.0 mm) in diameter
Particle size between 0.2 and 2 cm in diameter
Organic material in various stages of decomposition
Lime content
Reaction (pH)
Phosphorus content

Sample 1: Block 2, plot 719

Description	Unit	Result
Depth of sample	m	1.2
Clay (<2 µm)	0⁄0	28.8
Fine silt (2-20 µm)	0⁄0	23.1
Coarse silt (20-63 µm)	0⁄0	1.4
Fine sand (63-125 µm)	0⁄0	3.9
Fine medium sand	0⁄0	6.5
(125-250 µm)		
Coarse medium sand	0⁄0	10.2
(250-500 µm)		
Coarse sand (500-2000 µm)	0⁄0	26.1
Org. Mat.	0⁄0	1.8
Lime	0⁄0	0.0
pH-H ₂ O	-	5.4
Р		- 1

Sample 2: Block 2, plot 183

Description	Unit	Result
Depth of sample	m	1.2
Clay (<2 μm)	%	29.4
Fine silt (2-20 µm)	%	24.2
Coarse silt (20-63 µm)	0⁄0	3.9
Fine sand (63-125 µm)	%	4.4
Fine medium sand	%	6.5
(125-250 μm)		
Coarse medium sand	%	8.7
(250-500 μm)		
Coarse sand (500-2000 µm)	0⁄0	22.7
Org. Mat.	%	1.1
Lime	%	0.0
pH-H ₂ O	-	5.1
Р		- 1

Results noted as - 1: Amount not detectable

Results noted as - 1: Amount not detectable

Site description – Habinsaran

LOCATION

Province: North Sumatra District: Latitude (degrees and minutes): 02°17'29"N Longitude (degrees and minutes): 99°13'42E Altitude (m above sea level): 1315 m above sea level Managing office/institution: PT INTI Indorayon Utama, R&D Department Owner: do User(s): do

Distance to nearest office responsible for management of the trial (km): *30* Distance to nearest villages/towns (km): Number of inhabitants in the nearest villages/towns:

Type of area (e.g. research station, managed forest, etc.): Managed forest plantations, mainly *Eucalyptus* sp. for pulp

CLIMATE

Nearest weather station:

Name of the station: *Habinsaran base camp (2 km from trial site)* Latitude (degrees and minutes): *see above* Longitude (degrees and minutes): *see above* Altitude (m a.s.l.): *see above*

Climatic data1	Jan.	Feb.	Mar.	Apr.	May	June	July	Aug.	Sep.	Oct.	Nov.	Dec.	Year
Rainfall (mm)	101.1	79.9	163.1	205.9	165.1	84.8	117.2	147.9	217.5	157.8	173.4	150.9	1764.5
Temp. mean (°C)	20.3	20.7	20.6	21.2	21.7	21.4	20.8	20.9	20.6	20.6	20.8	20.4	20.8
Temp. mean max.2 (°C)	-	-	-	-	-	-	-	-	-	-	-	-	-
Temp. mean min.3 (°C)	-	-	-	-	-	-	-	-	-	-	-	-	-
Evapotranspiration4 (mm)	-	-	-	-	-	-	-	-	-	-	-	-	-

¹ Period of observations 1988-98

² Average of daily maximum temperatures

³ Average of daily minimum temperatures ⁴ Potential evapotranspiration (ETP) - Penman's formula

Rainy season:

Number/type of seasons: one two x Even

Period(s):

(specify months)	

Length of rainy season:

No. of intermediate days: (pre- and posthumid period of the growing season) No. of wet days: (growing season)

Number of dry months per year (< 50 mm rain/month): *None* Frost (number of days/year): *None* Prevailing winds (direction, period, speed): *W to SW very occasionally strong* Alternative weather station: Name of the station: Latitude (degrees and minutes): Longitude (degrees and minutes): Altitude (m a.s.l.):

Climatic data ¹	Jan.	Feb.	Mar.	Apr.	May	June	July	Aug.	Sep.	Oct.	Nov.	Dec.	Year
Rainfall (mm)													
Temp. mean (°C)													
Temp. mean max. ² (°C)													
Temp. mean min. ³ (°C)													
Evapotranspiration ⁴ (mm)													
 ¹ Period of observations ² Average of daily maximum temperatures ³ Average of daily minimum temperatures ⁴ Potential evapotranspiration (ETP) - Penman's formula Rainy season: Number/type of seasons: one two even/irregular													
Period(s): (s	specify	montł	ns)										
Length of rainy season: No. of intermediat No. of wet days:	e days:	(grow	(pre ing seas	- and p son)	oosthu	mid pe	riod of	the gro	owing s	eason)			
Number of dry months per Frost (number of days/year Prevailing winds (direction,	year (<): period	< 50 m l, speec	m rain l):	/mont	h):								
TOPOGRAPHY (slope) o	f trial s	ite											
x Flat/gentle (0-8%)		Interm	ediate (9-30%))	Steep	(>30%)					

GENERAL SOIL DESCRIPTION

Soil texture		Soil depth		Soil drainage/ n		Gravel content, topsoil		
1. Light/sandy		1. Shallow (< 50 cm)	Х	1. Well drained	Х	1. None (< 15 %)		
2. Medium/loamy	Х	2. Deep (50-100 cm)		2. Seasonal		2. Gravelly (15-35 %)	Х	
3. Heavy/clayey		3. Very deep (> 100 cm)		3. Permanent		3. Stony (> 35 %)		

Organic matter content		Reaction (pH)		Soil salinity		Groundwater		
1. Poor (< 2 % DM)		1. Acid (pH < 6.5)	Х	1. None	Х	1. Shallow (< 50 cm)		
2. Medium (2-5 % DM)	Х	2. Neutral (6.5-7.5)		2. Moderate		2. Deep (50 - 150 cm)		
3. Rich (> 5 %)		3. Alkaline (pH> 7.5)		3. High		3. Very deep (>150 cm)	Х	

Specify soil unit, soil association and phases (subdivisions of soil units) according to the Soil map of the world (FAO-Unesco 1971-1979), if known:

VEGETATION

Natural (original) vegetation type: Natural forest

Dominant natural (original) genera/species: Many

Land use history: Natural forest -> degraded forest/scrub -> eucalypts

RESULTS OF SOIL SAMPLE

Results of the laboratory analysis of the soil samples taken at the trial site.

The variables are:

Depth:	Soil sample depth
Clay:	Particle size less than 2 μ m (0.002 mm) in diameter
Fine silt	Particle size between 2 and 20 μ m (0.002 - 0.020 mm) in diameter
Coarse silt	Particle size between 20 and 63 μ m (0.020 - 0.063 mm) in diameter
Fine sand	Particle size between 63 and 125 µm (0.063 - 0.125 mm) in diameter
Fine medium sand	Particle size between 125 and 250 µm (0.125 - 0.250 mm) in diameter
Coarse medium sand	Particle size between 250 and 500 µm (0.250 - 0.500 mm) in diameter
Coarse sand	Particle size between 500 and 2000 µm (0.500 - 2.0 mm) in diameter
Gravel	Particle size between 0.2 and 2 cm in diameter
Org. mat.	Organic material in various stages of decomposition
Lime	Lime content
pH-H2O	Reaction (pH)
P	Phosphorus content

Sample 1: Block 2

Sample 2: Block 1

Description	Unit	Result
Depth of sample	m	1.2
Clay (<2 µm)	%	7.7
Fine silt (2-20 µm)	%	6.3
Coarse silt (20-63 µm)	%	4.9
Fine sand (63-125 µm)	0⁄0	9.3
Fine medium sand	%	18.4
(125-250 µm)		
Coarse medium sand	%	22.2
(250-500 μm)		
Coarse sand (500-2000 $\mu m)$	0⁄0	31.2
Org. Mat.	0⁄0	3.4
Lime	%	0.0
pH-H ₂ O	-	5.7
Р		- 1

Description	Unit	Result
Depth of sample	m	1.2
Clay (<2 µm)	0⁄0	23.1
Fine silt (2-20 µm)	0⁄0	8.2
Coarse silt (20-63 µm)	0⁄0	17.0
Fine sand (63-125 µm)	0⁄0	7.4
Fine medium sand	%	11.0
(125-250 µm)		
Coarse medium sand (250-500 μm)	0/0	14.9
Coarse sand (500-2000 µm)	0⁄0	18.4
Org. Mat.	0⁄0	7.5
Lime	0⁄0	0.0
pH-H ₂ O	-	5.7
Р		- 1

Results noted as - 1: Amount not detectable

Results noted as - 1: Amount not detectable

Annex 4. Plot data set

Aggregated data set at plot level. This annex describes the variables in the plot dataset, and displays the data. The plot data set has been prepared from the individual tree dataset and holds the following parameters.

PARAMETER NAME	EXPLANATION
SITE	Name of site
DATEEST	Establishment data of trial (MM/YY)
DATEASS	Date of assessment (MM/YY)
BLOCK	Block No.
PLOT	Plot No.
PLOTX	X- coordinate (see map)
PLOTY	Y-coordinate (see map)
SEEDLOT	Seedlot No.
PROVNAME	Name of provenance
SURV	Survival percentage (%)
DG	Diameter corresponding to mean basal area at breast height (cm)
GHA	Basal area (m2/ha)
GMEAN	Mean basal area per tree (m2)
HG	Height for tree with diameter corresponding to mean basal area (m)
MEANPILO	Mean pilodyn for plot
PILOCORR	Mean pilodyn reading adjusted for diameter effect
STEM	Average stemform
STEM1STEM9	Frequency of individual stemform scores 1 to 9 (%)
WHORLS	Average number of whorls
BRANCH	Average number of branches in whorl
DIABRA	Diameter of largest branch (cm)
FORK	Frequency of trees with one or more forks (%)
FO_POS	Average position of lower fork (m above ground)
FO_INDEX	Forking index (m ⁻¹)
FLOWER	Flowering and fruiting frequency (%)
FOXTAIL	Frequency of foxtails in plot (%)
KRAFT	Average Kraft index
KRAFT1 KRAFT5	Frequency of individual Kraft scores 1 to 5 (%)
CROWN	Average crown index
CROWN1CROWN5	Frequency of individual crown index scores 1 to 5 (%)
BR_INDEX	Branching index (cm)
INTNODE	Average distance between whorls (m)
BRARATIO	Ratio between branch diameter and DBH

DG- Diameter corresponding to mean basal area at breast height (1.3 m)

DG is calculated using the following formula:

where

D_i is the diameter at breast height of tree No. i (in cm);

n is the total number of trees in plot.

GHA - Basal area per hectare

Basal area in m² per hectare is calculated using the formula:

$$GHA = \frac{\sum_{i=1}^{n} \pi \times \left(\frac{Di}{2 \times 100}\right)^{2}}{n \times sp^{2}} \times 10000$$

where

D_i is the diameter at breast height of tree no. i (in cm);

n is the total number of trees in plot; and sp is the spacing in m;

HG- Height of tree with diameter corresponding to mean basal area

A linear regression per plot is prepared using the model:

$$Height = \beta + \alpha \times \ln(DBH1)$$

where

DBH1 is the diameter of stem (first stem if more than one stem) in cm; α is the slope of the regression line;

 β is the intercept with y-axis;

For each plot, α and β are estimated using PROC REG (SAS 1990).

HG for the plot is then calculated using the linear regression estimates (α and β) and plot DG (as previously calculated).

PILOCORR - Correction of pilodyn readings

Tree diameter (ring width) influences the pilodyn reading, i.e. trees with larger diameter (rings) will normally have larger pilodyn readings (deeper penetration of the pilodyn pin) than trees with smaller diameter, all other factors equal (ceteris paribus).

In order to adjust pilodyn readings for the variation in diameter, a correction factor has been introduced. By doing so, we are reducing the variance due to differences in individual tree size, and the provenances are in the subsequent analysis compared assuming that they have the same average tree size. In other words, we compare the level of the trait rather than the actual value.

The adjustment has been made using the GLM procedure in SAS (SAS 1990). The following model is applied:

. . .

. . .

PROC GLM; CLASS plot; MODEL pilo = plot DBH1; LSMEANS plot OUT=A;

Forking index

The forking index is calculated using the formula:

$$FO_INDEX = \frac{\sum_{i=1}^{n} FORK_i / FO_POS_i}{n}$$

where

FORKi is the number of forks observed on tree i FO_POS_i is position above ground of first fork (in m) on tree i; and

n is the total number of trees with forking data in the plot

Branching index

The branching index is calculated using the formula:

$$BR_INDEX = \frac{\sum_{i=1}^{n} BRANCH_i \times DIABRA_i}{n \times 10}$$

where

BRANCH, is the number of branches on tree i; DIABRA, is the branch diameter on tree i; and n is the total number of trees with branching data in the plot.

INTNODE - Average distance between whorls

The INTNODE parameter is calculated using the formula:

$$INTNODE = \frac{\sum_{i=1}^{n} (HEIGHT_i - 0.5) / WHORLS_i}{n}$$

where

HEIGHT_i is the height of tree i WHORLS_i is the number of whorls on tree i; and n is the total number of trees with observations on whorls in the plot.

Plot Data Set - Aek Nauli

м н н н н н н н н н н н н н н н н н н н	0 14.28571429 0 0 0	0 0 50 10 10 0 0 0.090909091 0 68.75		0000 000 0	00000	
S E M 6.4 5.285714286 5.285714286 5.285714286	6.6 6.666666667 4.285714286 3 5.66666667 7 7.363536364 3 6.583333333 4 6.533333333 4 7 7	7 5 5.90909090 3 6.88838889 3 6.888388889 3 6.888388889 12.5 6.166666667 4.909090909 6.333333333 6.333333333 5.2000090909	3.0.00.00.00 5.0666666667 1 5.8888888889 3 5.8888888899 5 7.25	6.571428571 6.5 6.8 7.1 6.8 57142857 6.4 61538462 5 6.4 61538462 5 5 5 5 33333333333333333333333333333	5 25 6 6 5 6.875 9 6.4545455 1 6.4545455 1 6.25	1 5.8888888889 4.8 4.8 5 5.888888889 4.8 4.4 6.88888888989 4.4 6.88888988989 4.4 6.846153946 6.333333333 6.3333333333 6.333333333 6.333333333 6.2 6.333333333 6.2 6.3 6.2 6.3 6.2 6.3 6.2 6.3 6.2 6.3 6.2 6.2 6.2 6.2 6.2 6.2 6.2 6.2 6.2 6.2 6.2 6.2 6.2 7.7777776 6.2
ς Χα∢τ⊢	0 3.72727272 0 3.083333333 0 3.083333333 14.668666665	0 3.818181818 92 3 0 1.7777777 0 1.7777777 25 6.25	0 3.22222222 0 3.222222222 0 3.77777778 5 4.25	0 3.428574429 0 3.16666661 3.5 0 3.5 0 12.5	50 3.5454545454 0 3.54545454545 0 3.126 0 3.126 0 4	0 3.4 0 3.5 0 3.5 0 3.5 0 3.4 0 2.888888888888888888888888888888888888
⊻⊻∢∟⊢∽	50 273 567 100 66,666666	182 388 7.6923076 0 0 3 0 3 0 6.5	50 444 100 444 22.222222	0 0 333 50	0 455 455 7.5 9.0909090	255 255 255 255 255 255 255 255
О Х К К Г Н 4	6667 2727 72.72727 75 16.66666 0 33.33333	1818 81.81818 4615 15.38461 0 12.5 0 0	50 3333 44.4444 0 2222 44.44444	50 33.3333 50 33.33333 0 0 0	0 4545 54.54545 37.5 3 1818 72.72727 0	30 3333 33.33333 100 33.33333 0 0 0 0 0 0 0 0 0 11.1111
⊻ ₩ < ₩ ⊢ ∞ ○	66667 16.6666 0 27.2727 133333 0	0 18.1818 23077 46.1538 77778 0 0	0 22222 33.3333 0 11111 22.2222 25	157143 42.8571 166667 40 0 0	0 0 45.4545 25 0 18.1818 0	20 66667 33.3333 0 20 50 50 50 50 66.6666
XK4FF4	3666667 16.066 0 8.3333 0 8.3333	0 0 30.765 5.5 5.5 6.625	0 22.222 0 11.111	0 7.1428 0 16.666 0 0 6.625 7	4.75 0 0 0	0 16.086 0 16.086 0 25 25 25 25 25
zoım-qı+ ^θ ννω	4 1 4 0 1 1 0 0 0 90 90	1108011011	048-04	⊈∞ ⊏≰ο≅≻ຬ∞	4 E @ E 4 ©	5 r c o c o c o c o c o c o c o c o c o c
20 10044196666 20 106874	6 6 6 6 6 6 6 6 6 6 6 6 6 6 6 6 6 6 6	66666666666666666666666666666666666666	5 6 6 6 9 9 9 5 4 6 7 6 9 9	5 5 5 5 5 5 5 5 5 5 5 5 5 5 5 5 5 5 5	66666 - 6666 - 4787 - 47877 - 47787 - 4777 - 4777 - 4777 - 4777 - 4777 - 4777 - 4777 -	φ φ
LOXF<	50 53.3333333 53.63636364 50 50 50	2.727273846 2.72727273 25	50 50 50 50 50 50 50 50 50 50 50 50 50 5	8.57142857	33.63636364	100 50 53.33333333 33.33333333 70 70 80 80 83.33333333
ΓΟΚΥ	0.4 0.2727273 0.5 0 0	0.909090909	0.5 0.33333333 (0.25 0.75	0.090909090	0.3636364 (0.375 0.454545455 (0.454545455 (0.5 100	0.285714286 0.285714286 0 0.6 0.6 0.6 0.75 0.4444444 3
$\Box - \triangleleft \square \square \prec$	5.1 4.67 5.333333333 4.75 8.3	6.245454545 5.51111111 3.414285714 3.414285714	3.875 4.87777778 3.2 5.822222222 5.75 5.75	6.7 5.65 2.75	6.32 7.1 5.336363636 3.6 0	5.866666667 4.95 3.133333333 3.133333333 3.133333333 3.133333333
n r < z ∪ ī	7 4.66666667 2 5.090909091 3 8.916666667 9 4.3333333333333	7.272727373 12.909090 5.7777778 5.77777778	5 4.5 4 4.88888889 5 7.22222222	5 5.928571429 5 5.33333333 8 4.3 4.3	5 6.636363636 5 4.75 5 5.818181818 5 5.818181818 5 5	4.6 4.6 4.6 4.6 4.7 5 5 4.7 5 5 4.7 5 5 5 4.7 5 5 5 5 5 3.875
SIOK⊣0	5 7 7 3 10.1666666 1 8.18181818 7 7.58333333 5 8 8.33333333 9 8 8.33333333	10.0000000 4 4.818181819 7 4.777777 6 7 7 7 7 7 7 7 7 7 7 7 7 7	5 16.7 1 7.4444444 2 8.7777777	4 6.78571428 5 6.1 5 5 5.1 5.1 5.3 3 5 5 5 5 3 3 5 5 5 5 5 5 5 5 5 5 5 5	3 6 11.4545454 5 7.622 2 8.454545454 5 16 5 16 5 16 5 16 5 16 5 16 5 16 5 16	7.16666666 7.166666666 3 3 3 3 3 3 3 3 3 3 3 3 3 3 3
7 4 8 2 2 2 2 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1	6 22.6923076 6 22.6923076 4 4 1 1 20.8333333 7 19.4166666 7 19.4166666 1 16: 1 16:	1 19.590990 5 19.1363636 6 19.4166666 6 2187 6 18.7 6 18.7 7 22.4615384 1 22.4615384 7 20.2727272 7 20.2727272	4 17.3030303 1 23.62 3 18.1111111 3 18.1666666 3 18.1666666 1 17.7	6 19.6428571 7 16.1666666 5 17.687 5 23.6071428 5 23.6071428 5 23.6071428 6 23.602 6 23.562 6 21.66 6 20.4285714	1 20. 4 21.3636363 5 23.2 5 23.2 5 18.62 7 22.	7 2 21. 8 9 9.9686666 9 9 9.9686666 6 9 9 9.1 1 21.66666666 6 9 9 9 9.2 1 1 18.8333333 2 2 2 7 1 9.8571428 9 2 2 2 2 2 4 4 4 4 4 4 4 4 4 4 4 4 4 4
G G C C C C C C C C C C C C C C C C C C	57 0.13767128 68 0.67837078 21 0.3036542 62 0.31925355 29 0.3511522 28 0.3511522 28 0.3614522 28 0.36438 76 0.15475192 76 0.15475192	43 0.47372757 95 0.278184 96 0.2781858 48 0.38022677 15 0.32646976 44 0.49914857 35 0.4020712 35 0.4020712 36 0.12742635 37 0.62196435 37 0.62196435 37 0.62196435 38 0.127742635 39 0.127742635 30 0.127742635 30 0.127742635 31 0.4020712 32 0.127742635 33 0.4020712 34 0.40976 35 0.4020712 35 0.402075 35 0.40207	28 0.317/00/ 68 0.1591770 94 0.2325266 43 0.02351564 94 0.34612914 43 0.22243251	99 0.33083676 26 0.16927035 75 0.083955 67 0.282745 05 0.119102 63 0.4119102 63 0.4110582 53 0.37146528 12 0.2915975	14 0.20005307 29 0.4442232 15 0.210630 75 0.19741 57 0.04572930	86 0.3302168 84 0.12640728 43 0.032663033 43 0.032663033 43 0.03266675 53 0.1514896- 53 0.12296495 53 0.22954956 60 0.30912981 60 0.30912981 61 0.30557025 44 0.3055781 44 0.3055781 53 0.0055781 53 0.0055781 53 0.0055886- 53 0.005896- 53 0.005896- 54
G G G E E E A A A A A A A A A A A A A A	1.25 0.0275342 1.25 0.0521823 8.25 0.0303654 8.75 0.0365422 8.75 0.0319229 75 0.02814045 8.75 0.02818	8.75 0.0430661 8.75 0.02528 5.25 0.0081306 5.0 0.0475283 75 0.0272058 75 0.0335059 37.5 0.0338050 37.5 0.0338050	8.75 0.0298924 25 0.0397942 6.25 0.0258362 6.25 0.0235156 6.25 0.0384587 25 0.0556081	87.5 0.0236311 37.5 0.0282117 37.5 0.008395 3.75 0.0186499 3.75 0.0186499 1.25 0.02302 1.25 0.0316198 50 0.0560562 50 0.0560562 3.75 0.01665742 3.75 0.0416571	1.25 0.0400106 8.75 0.0403839 50 0.0263291 8.75 0.0361 25 0.0493528 6.25 0.0457293	22.5 0.0330216 23.75 0.0190561 25.5 0.0276946 25.9 0.02659816 25.5 0.02659813 27.5 0.026797 27.5 0.03008713 27.5 0.03008713 27.5 0.03008715 27.5 0.0296713 27.5 0.0296713 27.5 0.0296713 27.5 0.029615 27.5 0.029615 27.5 0.01763951 22.5 0.01763951 23.5 0.0176355 23.5 0.0176555 23.5 0.01765555 23.5 0.01765555 23.5 0.0176555555555555555555555555555555555555
$\omega \supset \pi > _{\omega} _{4}$	~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~		2 2 2	0000 4	~ ~ ~	4 − 0 0 4 − 0
Н Е Е С В 9.657142857 11.442857142857 11.76666667	9.95 14.96363636 12.45 10.46666667 8.345454545 7.3 7.3 9.266666667 7.3	9.5545455 7.245454545 4.422222222 11.6125 10.09 12.66 11.83636364 9.93333333 14.28	8.77 11.35 7.2625 6.8 9.566666667 9.975	8.335714286 7.08333333 4.654545455 9.085714286 11.16923077 13.71428571 13.71428571 11.82307692 11.91666667	10.425 11.9 7.9375 8.154545455 10.95 9.5	8.87 6.585714286 9.3 8.77777777 9.3 8.7777777 10.14 10.14 10.14 10.2333333 12.3333333 12.3333333 12.3333333 12.3333333 12.3333333 12.3333333 12.3333333 12.3333333 12.3333333 12.33333 12.333333 12.333333 12.333333 12.333333 12.333333 12.333333 12.333333 12.333333 12.333333 12.333333 12.333333 12.333333 12.333333 12.3333333 12.333333 12.333333 12.333333 12.333333 12.3333333 12.3333333 12.333333 12.333333 12.33333333 12.333333333 12.3333333333
D B H 14.1733333 5 18.88571429 5 18.88571429 5 22.3	17.48 25.39230769 18.99 20.46666667 19.9 19.9 18.26666667 18.26666667 22.556666667 22.556666667	23.21818182 16.41818182 5 16.41818182 5 7.96556556 5 7.96556556 5 24.3625 5 18.44166667 5 21.89230769 5 20.15833333 5 20.15833333 5 20.15875 5 20.15975 5 20.15875 5 20.15975 5 20.159755 5 20.159755 5 20.159755 5 20.159755555555555555555555555555555555555	18./0305030 22.125 17.7444444 17.3 21.53333333 21.53333333 22.5 22.5	16.86428571 18.71666667 9.47 9.47 9.47 9.47 9.47 9.47 17.16666667 17.406666667 17.261538466 26.4125 18.55384615 218.55384615 22.64285714	22.3 22.5 18.1125 20.60909091 24.525 24.525 24.525 5 24.525 5 24.525 5 24.525 5 24.525 5 24.525 5 24.525 5 24.525 5 24.525 5 24.525 5 24.525 5 22.52 5 22.55 5 5 22.55 5 5 22.55 5 5 5	19.36 14.25714286 18.7 19.3 19.2 19.2 19.2 19.2 19.2 10.2 10.2 10.3 11.344444 11.34444444
IF>FM 10000	000000000			000000000		
с - о - х - х - б - р - р - х - 2 - 2 - р - 2 - 2 - 2 - 2 - 2 - 2 - 2	× & o 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0	5 2 2 3 3 4 5 5 3 3 4 5 5 3 3 4 5 5 3 3 4 5 5 3 3 5 5 5 3 3 5 5 5 5	04 05 05 05 05 05 05 05 05 05 05 05 05 05	2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2	5 42 4 5 24 4 5 56 4 5 5 5 5 5 5 5 5 5 5 5 5 5 5 5 5 5 5 5	 8.45 8.45 8.45 8.67 9.60 9.61 9.65 <l< td=""></l<>
000 000 000 000 000 00 00 00 00 00 00 0		0.22 30 30 30 30 30 30 30 30 30 30 30 30 30	20.26		0.26	0.20 0.20
σ-10+× ₀ ∞υ _ω σ-10+×04466	~ @ @ @ 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0	vor55ov5ro 4441500500	×65004	8 4 8 4 8 4 6 4 6 6 1 0 1 4 8 6 4 6 6	o400≻0 47700€	© < < < < < < < < < < < < < < < < < < <
□ < ⊢ □ < ∪ < < < < < < < < < < < < < < < < <	999 2 999 2 1 1 1 1 1 1 1 1 1 1 1 1 1 1	9999 9999 9999 9999 9999 9999 9999 9999 9999	999 999 999 999 999 5 7 7 7 7 7 7 8 9 9 9 9 9 9 9 9 9 9 9 9 9	9999 5 9999 5 9999 7 9999 8 9999 8 9999 9 9999 7 9999 9 9999 9 9999 9	000 000 000 000 000 000 000 000 000 00	9999 6 9999 6 9999 7 9999 9 9999 8 9999 8 900 8 9000 8 900 8 900 8 9000 8 9000 8 9000 8 9000 8 9000 8 9000 8 9000 8 9000 8 90000 8 9000 8 9000 8 9000 8 9000 8
а – – – – – – – – – – – – – – – – – – –	1192 1192 1192 1192 1192	1192 1192 1192 1192 1192 1192	1192 1192 1192 1192	1192 1192 1192 1192 1192 1192	1192 1192 1192 1192	1192 1192 1192 1192 1192 1192 1192 1192
s AekNaufi AekNaufi AekNaufi AekNaufi	AekNauli AekNauli AekNauli AekNauli AekNauli AekNauli AekNauli AekNauli	AekNauli AekNauli AekNauli AekNauli AekNauli AekNauli AekNauli AekNauli	AekNaul AekNaul AekNaul AekNaul AekNaul	AekNaufi AekNaufi AekNaufi AekNaufi AekNaufi AekNaufi AekNaufi AekNaufi AekNaufi	AekNauli AekNauli AekNauli AekNauli AekNauli AekNauli	AekNauri AekNauri AekNauri AekNauri AekNauri AekNauri AekNauri AekNauri AekNauri AekNauri AekNauri AekNauri AekNauri
R R R R R R R R R R R R R R R R R R R	1 714 OOCARRA(HOND) 715 TECUNUMANII(RAP) 1 716 CARIBAEA(GUA) 718 COTOMINES 1 719 NONGRATING 1 720 NONGRATING 1 721 LANGHAMH 1 722 DOISUTHEP	1 728 BODANA(NB) 1 728 BUDANA(NB) 1 728 SHANGSI 2 73 SHANGSI 2 118 PATULA(ZBW) 2 138 MERCUSI(INI) 2 713 OCCARPA(MALPASC 2 713 COCARPA(IAPOND) 2 715 TECUNUMANII(RAP) 2 715 TECUNUMANII(RAP) 2 715 TECUNUMANII(RAP)	2 716 CATRBEAG(GUA) 718 CATRBEAG(GUA) 2 719 NAMNOW 2 720 NONGKRATING 2 721 LANGHANH 2 722 DOISUTHEP	2 728 BODANA(AB) 2 728 ANUSBAN 2 728 SHANGSI 3 118 PATULA(ZBW) 3 133 REACUSI(IND) 3 713 COCARPA(MALPSC) 3 713 COCARPA(MALPSC) 3 715 TECUNUMANI(RPS) 3 715 TECUNUMANI(RPS) 3 715 TECUNUMANI(RPS)	3 716 CATBREAG(GUA) 718 COTOMNES 719 NAMNOW 720 NONGKRATING 721 LANGHANH 722 DOISUTHEP 7 73	7 22 42 42 42 42 42 42 42 42 42 42 42 42

	_	ш ;	> ц		14	13	13	5	10	;	12	12	5	÷	12	13	5	;	12	. 16	17	7	10	15	12	13	15	. 16	44	15	10	14	6	12	2	00	s ç	<u></u> 9	,	ი ი	8	б 	4	6	2	10	;	00
		2	> I	. ⊲	1.553228	6271493	8747775	0410243	.3848372	1.540269	1.942926	2133003	1.263807	7484063	3863756	.1407882	0.472736	.1465639	4962328	7.296717	8.059819	0.523396	4.269419	.071290	6.73713	.1287646	1.827566	.8450594	5864755	4.343172	.3742166	7.990624	6660842	7778152	.0129968		8.534837	0.10000	10000000	0981996	7.748131	5925630	0.621197	8043783	8228676	0.281643	.6222154	.2318336
					373 9	317 74	691 81	877 64	366 45	029 35	673 18	219 81	743 10	296 75	468 34	353 47	496 15	419 73	859 17	867 14	245 11	899 20	454 16	809 49	016 29	003 98	405 6	434 58	537 5	238 11	131 80	914 9	834 41	514 20	794 98		235 17	90/ 19 206 16		877 75	494 17	443 58	915 11	448 73	918 14	349 10	344 32	979 26
₽ – ⊐	0	00	2 0	<u> </u>	23.1835	20.9449	20.9377	22.0083	21.2028	23.7175	17.300	21.0353	19.0163	19.0685	16.916	19.0543	20.2527	18.6617	17.8136	22.7280	18.0769	22.9017	20.4377	20.592	21.9196	17.281	24.1041	17.8581	21.6727	18.5469	18.2918	19.2427	16.0761	16.2697	22.8429		20.2802	1907.42		20.8083	21.9054	23.0585	22.2257	19.5052	22.228	20.0170	20.9276	19.0733
			Е	<u>ں</u> ב	9.48043514	9.62338276	11.7452148	12.2214009	11.1016394	14.9514421	10.3617019	10.9376228	8.46895188	7.45813892	1.37241538	9.26666657	9.5865563	7.45442701	5.37640071	11.731436	10.0710771	13.2769335	12.0181367	10.7444009	14.3918021	9.10580383	11.5408346	7.48687173	6.8	9.79496098	11.2711021	8.52270046	7.13877451	5.44054167	9.36428316		12.5719153	18/0188.01	110440.21	10.4173178	12.0313527	8.041223	8.2183125	11.2037622	9.5	8.85882207	7.07233145	9.33068169
			6	ט נ	15.0496512	18.5488544	19.2844127	24.5983739	18.7199359	25.7708989	19.6588148	21.5657908	20.1566817	17.493737	18.9203594	25.6227113	23.4118463	17.9406344	10.1725666	24.5948419	18.6079508	22.1060382	20.6504035	16.4407522	22.24285	19.1727553	22.5049439	18.1335478	17.3	22.1240945	26.6033832	17.3424541	18.9488346	10.3370692	15.4889853	17.4117776	20.0607731	20.7103078	72 0756690	22.5660364	22.6710669	18.3057027	21.4393436	25.0624719	24.1248834	20.5006341	15.1601922	18.7709883
			C	οo								-	-			-	-	- -	-								-	-		-	-	-	-	-							-	-	-	-	-	-	-	-
			, ,									-	.	.	- ·	~	.	.	.								15	.	- ·	-	.	.	.	-							.	-	-	-	.	5	.	.
			ц	- LL								.	-	. -	-	-	.		-								-	.	~	-	-	-	.	-							.	-	-	-	-	.	.	-
			W	ΣLL								2	1.27272727	~	- -	4	2.81818182	1.27272727	1.42857143								2.5	2	m	7	°	1.35714286	1.83333333	1.4							2.09090909	1.125	2.727273	2	-	2	1.42857143	~
	>	⊢ c	2 Ц	чш	0.0878911	0.13432887	0.16842811	0.23054769	0.13070833	0.38939845	0.26199781	0.19491192	0.13256353	0.09089809	0.1650546	0.22627578	0.19698249	0.0957555	0.02799397	0.26513409	0.14167178	0.22211822	0.1971233	0.1177711	0.26706342	0.12845947	0.22257924	0.0941521	0.08044525	0.18294908	0.28934718	0.10079036	0.0999986	0.02992005	0.09409248		0.19776167	0.35312499	0.2250052	0.21628281	0.23268846	0.10546661	0.1448132	0.26569576	0.21344929	0.14440557	0.06710856	0.1259128
		۲	- 0	0 ₪	0		0					0	0	0	0	6.25	6.25	0	0				0	0			0	0	0	0	0	6.25	0	0	0		0	c	>		0	6.25	6.25	12.5	0	6.25	0	0
ш	0	10	LC	იი								2.3	3.333333333	4.18333333			3.25	1.5222222									4.9	4.45		3.6	ę	4.6									4.6	3.23333333	2.725	5.2		2.6	1.9	
vĸo	N	z	LC) H								80.6122858	68.5336114	70.762255	87.1916843	84.4721373	63.879367	68.6977716	59.3976578								85.4976438	80.9426933	82.3529412	84.6786501	88.3232387	72.2203313	77.0685096	77.1468507							79.0265247	75.035373	80.3024771	85.5024846	81.0526316	72.7844121	73.9469646	68.5215593
– z	F	z		ш								1.0025463	1.1007684	1.05498764	1.12083333	1.05648148	0.85177386	1.51555556	0.97441799								0.68227606	1.14129464	1.26	1.05494468	0.90360119	1.24653912	1.05888889	0.93268254							1.51720822	1.28619048	0.94277056	0.77362013		1.17636111	1.07478175	0.76106061
шО	ı —	z	ш с	ч×								5.47420635	2.69888017	2.3940734			2.88986355	7.91418357									2.56008772	1.70960366		3.05660934	5.20151515	1.86956522									3.4334507	2.87426901	15.6230447	4.11538462		4.03188859	4.16769231	
<u> </u>	Ľ	×۲		- 0		0			6.25	12.5	18.75	0.23762695	0.22497786	0.30520457	0.21823467	0.31739985	0.26182314	0.31467897	0.54048238		0	18.75				6.25	0.17830331	0.28394681	0.1849711	0.27461816	0.17586116	0.35875336	0.30721651	0.32090803					c	0	0.29110162	0.38040502	0.25945332	0.14699179		0.34819648	0.30839714	0.16179295
	S	н L	μZ	≦ 0	0	0	0		0	0	0	0	60606060.6	0	0	0	0	0	0		0	0	0	0		0	0	0	0	0	50	7.14285714	0	20	0		0	c		>	0	0	0	0		0	0	0
	S	н .	⊔≥	≥œ	6.66666667	0	14.2857143		60	25	0	0	27.27273	16.6666667	0	0	27.27273	18.1818182	11.111111		0	0	60606060.6	16.6666667		0	25	11.1111111	0	22.222222	0	21.4285714	16.6666667	10	42.8571429		15.3846154	11 666667	1000000.14	þ	0	25	60606060.6	25		11.111111	0	33.3333333
	s	н L	⊔≥	7	40	0	14.2857143		20	41.6666667	14.2857143	33.3333333	54.5454545	41.6666667	100	33.3333333	54.5454545	9.09090909	66.6666667		10	50	18.1818182	50		9.09090909	0	33.3333333	0	22.222222	0	35.7142857	33.3333333	50	35.7142857		53.8461538	11 666667	41.0000001	8	60	50	45.4545455	25		11.111111	20	0
	S	н L	⊔ ≥	<u>8</u> 9	40	0	0		0	25	14.2857143	16.6666667	60606060.6	33.3333333	0	66.6666667	60606060.6	36.3636364	22.222222		0	33.3333333	60606060.6	0		60606060.6	25	22.222222	0	22.222222	25	21.4285714	33.3333333	10	0		15.3846154	c		>	10	12.5	36.3636364	25		33.3333333	20	0
	S	⊢ L	⊔≥	∑ Ω	13.3333333	0	42.8571429		0	0	14.2857143	33.3333333	0	0	0	0	60606060.6	27.27273	0		0	8.333333333	27.27273	16.6666667		60606060.6	0	11.1111111	0	11.111111	25	0	16.6666667	0	14.2857143		0		16 666667	0000000	10	12.5	0	0		44.444444	20	0
	S	⊢ι	υZ	<u>8</u> 4	0	14.2857143	14.2857143	6.25	0	0	28.5714286	16.6666667	0	8.33333333	0	0	0	0	0	0	0	0	60606060.6	16.6666667	0	9.09090909	0	0	0	11.111111	ō	7.14285714	0	10	0		7.69230769	0	0.0000000	0 0	10	0	60606060.6	25		0	20	66.666667
	S	н I	⊔≥	≦ ო	0	28.5714286	14.2857143	0	0	8.33333333	14.2857143	0	0	0	0	0	0	9.09090909	0	0	20	8.33333333	18.1818182	0	0	36.3636364	50	11.1111111	0	0	0	0	0	0	7.14285714		7.69230769 î	D	000000000000	0	10	0	0	0		0	0	0
	S	⊢ L	⊔≥	2 2	0	42.8571429	0	0	20	0	0	ō	Ō	0	0	0	0	0	0	0	60	0	ō	0	6.25	27.2727273	0	11.1111111	100	11.1111111	0	7.14285714	0	0	0		0 1	C.71	c	0 0	0	0	0	0		0	20	0

Plot Data Set - Aek Nauli

a,
ar
S.
<u>.</u>
al
Т
_
Ū
S
а
at
õ
)t
Ę

⊻╙⋖╙⊢	7			0		;	20.	31.25	37.5	18.75	1.2857143 5555557	/000000.1	b		0		3.6666667	3.0769231	20	3.1818182	33333333 67 E	2.20	8		0	0		25	0.6666667	3.1818182	20	0	1.7692308				c	0	3636364	20	5.3846154 20
⊻╙∢⊩⊢	-			9			0 566667 26	0	0	6.25	1 0 14	7.25	24		9		0 16	0 25	0	727273 18	xi D C	957143			5.9375	7		6.25	0 16	090909 15	0 366667 13	1 000000	692308 3C			0	a	6.5	0 36	0 0	20 II
		n 0	-	80	8	0	6 5 6.66	9	7	9	- c	0 0 0 1 4	o 4	2	0	е	0	9	5	1 27.2	N 0	4 14 2	. 6	2	9	2	Nσ	2	2	60 ⁻⁶	0 3 668	0.00 3 0.00	4 30.7	.	. -	4 (თ . ო	, œ	6	0.	~ ∞
zoimior	ים :	5 C	12 1	12	10	15	15 1	16 1	ø	16 1	4 (4 (<u>ν</u> α	15 1	14	12	15 1	12	13	15 1	÷ ;	- 21 0	o 4	12	15 1	16 1	9	22 C	16 1	12 1	7	10 10 10	<u> </u>	13 1	11	15	4	<u></u>	, <u>6</u>	£	10	13
ZOI⊢O⊢∢	4	16	16	16	16	16	53 16 50 16	5 16	5 16	5 16	4 16 15	0 4 0	16	4	16	16	37 15	5 16	37 16	55 16	53 16 16	16	16	16	16	16	16	5 16	3 16	5 16	50 16 16	0 16	1 16	15	16	16	13	16	5 16	0 16	16 16 10 16
∟ 0 × ⊢ < −	_						53.33333 6	81.2	2	68.7	71.428571	•					66.666666	61.538461	66.666666	45.454545	58.333355	35.714285						62	33.33333	45.454545	23 33333	10.10	23.076923						54.545454	9	84.615384
шОш	¥						0.26666667 0.3333333333	0.1875	0.125	0	0.42857143	67.0					0.75	0.53846154	0.333333333	0.18181818	0.41666667	0.35714286						0.1875	0.41666667	0.54545455	0 13333333	0.66666667	0.15384615						0.63636364	0.3	0.53846154
□ – < ∞ m	A						3.74 5.31538462	3.05	3.4625	3.34	3.69285714	701010107					4.55	5.03076923	5.69333333	3.29090909	4.16666667 2.05	3 43846154						3.333333333	3.56666667	3.9	3.52 3.5066667	3.86666667	2.28333333						4.15454545	3.52	3.16153840 2.74
m ừ ∢ Z U	т						7.8666667 8.4	6.5	7	7.1875	8.21428571 7 nf	C7. 1					8.91666667	8.84615385	8.26666667	7.54545455	6.58333333 F 07E	0.070 4.64285714						7.125	6.91666667	2	5.9 6	0 12.6666667	6.07692308						5.45454545	8.9	5.07692308 6.5
≥ток⊣	S					:	12 7.86666667	10.25	9.5	9.0625	7.57142857	1000001 +. 1					8.5	6	11.0666667	7.81818182	10.5833333 6 276	6.85714286						10	11.4166667	12.0909091	10.5	6.333333333	5.38461538						11.4545455	7.6	12 9.6
∑ш∢∠∟–	0	z1.9090909 19.3333333	22.35	23.2727273	23.8	22.9	21.5333333 19.3333333	21.3125	23.1875	21.75	20.8571429	020.12	19.9	24.1071429	21.625	24.3	21.75	22.1538462	21.3333333	20.125	21.25	21.2142857	17.944444	20.6785714	24.625	21.5555556	23.75	21.65625	22.6666667	21.5	22.35 วว 633333	22.3333333	20.5	20.1	19.9285714	23.35/1429	23.0384615	23.5	18.5	20.65	20.3846154 20.75
⊂ v ט	M	25721771	27089071	38672543	.29601864	0.551023	43726414	47439543	20823943	.40152279	30488936	00154043	45391579	49003664	.46189471	.54688621	.33853914	28976986	33948043	.18178836	44/52/14 0 10106	12488221	11794986	39065086	49234193	39090621	.3386/664 52311836	34077686	34197821	.32146243	28515143	04098757	05272929	.19507007	46217679	35979586	107072020	0.4273885	.24706393	20825279	417/3679
0 ≥ ш ∢	N 001201	978598 0.	257423 0.	322712 0.	960186 0.	673487	2915094 0. 2512699 0.	964971 0.	602993 0.	509517 0.	2177781 0.	144255 0	026105 0.	500262 0.	849123 0.	645908 0.	1282116 0.	228999 0.	263203 0.	652621 0.	1/29393 U.	892016 0	982915 0.	604339 0.	077137 0.	909062 0	822305 0. 1435932 0	129855 0.	849818 0.	922386 0.	2851514 0.	366252 0.	040561 0	773364 0.	081179 0.	1256997 0.	1532/04 0.	561571 0	246036 0.	082528 0.	1321336 U. 579547 0.
ω – «	V 61.0F	81.25 0.0	75 0.02	75 0.03	62.5 0.02	93.75 0.03	93.75 0.02 93.75 0.02	100 0.02	50 0.02	100 0.02	87.5 0.02		93.75 0.03	100 0.03	75 0.03	93.75 0.03	80 0.0	81.25 0.02	100 0.02	68.75 0.0'	20.0 cl	87.5 0.00	75 0.00	93.75 0.02	100 0.03	62.5 0.03	20.0 67	100 0.02	75 0.02	68.75 0.02	62.5 0.02 03.75 0.02	33.73 0.00	93.75 0.0	133333 0.01	93.75 0.03	93.75 0.0	76 0.00	75 0.03	68.75 0.02	62.5 0.02	81.25 U.(62.5 0.02
тш— 0 т	T	692.308 10.43	363636	14.975	14.7375	14.57	10.15 333333	11.9125	714286	1.48125	454545	0.9 222223	928571	416667	13.13	846154	9.76	230769	9.8	363636	166667 6 025	714286	88889	166667	13.05	142857	13.325 77778	333333	9.375	44444	10.03 076923	010923 666667	714286	545455 73.30	909091	142857	222222	15.225	777778	8.81	090909 10.9625
	20 0 0000	6923 9.37 12308	6.175 12.2	6667	18.93	13333	9.41 17.08 9.41	8.975	7.975 11.3	51875 1	85714 8.45	11 8 8 58	18.9 12.5	20.6 16.9	6667	20.78 15.4	18.05	3846 9.39	16.24	27273 7.56	123333 11.1	8571 5.55	6667 7.08	11.8	9375	21.7 16.8	18 1975 147	15.75 9.29	17.75	9091 11.2	18.3 15.88.9.03	13.333 6.46	5385 3.90	3636 10.3	3333 12.4	17.35 12.9	17 0 11.1	11.2 17.2	1818 9.37	15.82	17692 1u.1 17.8 ·
	- 100	5 14.865	0 7	5 19.491	6	3 19.373	5 18.146 5	0 -	6 1	3 17.5	5 15.428	0.010		6	3 19.616	0	6	3 15.515	6	5 13.172	5 Z1.330	5 8.8145	3 10.766	3 17.553	3 17.0	6	, , ,	1	6	3 17.890		5 12.533	5 6.4461	5 14.336	5 18.633	0	5 20.81t	5 20.083	3 16.518	0	ດ 18.9.5 ດ
ін х ш о	.		0	0	0	0	 -	0	0	0			0	0	0	0	0	0	0	- ·			0	0	0	0	 -	0	0	0	0 0	0	0	0	0	- · - ·	 -	, - , 0	0	0 0	 - 0
	×	12	4	5	9	~ '	<u>ა</u> თ	18	16	10	~ ç	2 09	65	35	48	36	44	40	55	8	200	404	13	22	30	28	55 GE	9	24	20	38		4	78	72	105	6.6	45	20	06	84 98
	× ۲	0 0 4 8	5 60	5 75	5 90	5 105	5 12C	5 126	5 112	22 70	5 5 5 5 5 5 5 5 5 5 5 5 5 5 5 5 5 5 5		5 C	5 45	5 16	5 28	5 20	5 24	55	200	0 4 2 7	5 4 7 4	1.0	5 10	5 18	10 1	0 4 1 0	i u	2	5	ю ч (ч	0 7 7	ι Ω	5 18	5 24	ר ה ה ה	οu Γ	,	5 10	ب ب م	17 IS
	~ ~	42.2	56.2	56.2	56.2	56.2	56.2	42.2	42.2	42.2	42.2	2000	20.2	2.2	12.2	0.2	6.2	2.2	6.2	9.0			20.2	6.2	2.2	30.2	2.02	2.2	12.2	2.2	12.2	12.2	30.2	20.2	12.2	42.2	20.2	42.2	30.2	42.2	30.2
⊾ ⊣ 0 ⊢ ×	2	0.25	2.25	0.25	0.25	2.25	6.25 12.25	12.25	6.25	0.25	2.25	30.0	0.25	0.25	2.25	2.25	2.25	2.25	0.25	0.25	0.20 30 0	0.25	20.25	12.25	6.25	12.25	6.25 6.25	20.25	12.25	12.25	6.25 20.25	20.25	20.25	0.25	0.25	2.29	2.25	6.25	0.25	0.25	0.25
	> {	4 6 7 7 7	4 15	5 15	6 15	7 15	9 15 15	9 14	8 14	5 14	4 4 4 4	2 ₹	1 10 20 13	59	4	4 7	4 5	4 6	5 5	5 10	0 v 9 Q		, t , c	2 5	3 6	2	ი ო ი ო	1 6	2 4	2 6	ω , 4 π	 - 4	1	6 3	6		 	1 m	5 2	6 1	6 Z
	s g	666	666	666	666	666	666 666	666	666	666	666	000	666	666	666	666	666	666	666	666	666	666	666	666	666	666	666 000	666	666	666	666	666	666	666	666	666	666	666	666	666	666
О∢⊢шш⊘	+ ,	1092	1092	1092	1092	1092	1092	1092	1092	1092	1092	1102	1192	1192	1192	1192	1192	1192	1192	1192	1192	1192	1192	1192	1192	1192	1192	1192	1192	1192	1192	1192	1192	1092	1092	1092	1092	1092	1092	1092	1092
ω – Γ	Ш	Habinsar Habinsar	Habinsar	Habinsar	Habinsar	Habinsar	Habinsar Habinsar	Habinsar	Habinsar	Habinsar	Habinsar	Habinear	Habinsar	Habinsar	Habinsar	Habinsar	Habinsar	Habinsar	Habinsar	Habinsar	Habinsar	Habinsar	Habinsar	Habinsar	Habinsar	Habinsar	Habinsar Hahinsar	Habinsar	Habinsar	Habinsar	Habinsar	Habinsar	Habinsar	Habinsar	Habinsar	Habinsar	Habinsar	Habinsar	Habinsar	Habinsar	Habinsar Habinsar
ω □ □ Ο Ο 0 □ □ Ο Ο − Ο		1 118 PA IULA(2BW) 1 183 MERKUSII(IND)	1 712 OOCARPA(MALPASO)	1 713 TECUNUMANII(MPR)	1 714 OOCARPA(HOND)	1 715 TECUNUMANII(RAP)	1 719 NAMNOW 1 720 NONGKRATING	1 721 LANGHANH	1 722 DOISUTHEP	1 725 BODANA(A8)	1 726 AUNGBAN		2 183 MERKUSII(IND)	2 713 TECUNUMANII(MPR)	2 714 OOCARPA(HOND)	2 715 TECUNUMANII(RAP)	2 719 NAMNOW	2 720 NONGKRATING	2 721 LANGHANH	2 722 DOISUTHEP	2 /25 BOUANA(A8)	2 727 SHANGSI	3 118 PATULA(ZBW)	3 183 MERKUSII(IND)	3 712 OOCARPA(MALPASO)	3 713 TECUNUMANII(MPR)	3 715 TECLINLIMANII(RAP)	3 719 NAMNOW	3 720 NONGKRATING	3 721 LANGHANH	3 722 DOISUTHEP 3 725 BODANA(A8)	3 726 AUNGBAN	3 727 SHANGSI	4 118 PATULA(ZBW)	4 183 MERKUSII(IND)	4 712 OUCARPA(MALPASO)	4 /13 LECUNUMANII(MPR)	4 715 TECUNUMANII(RAP)	4 719 NAMNOW	4 720 NONGKRATING	4 721 LANGHANH 4 722 DOISUTHEP

>	I∢	8.6391009	10.525896	26.161868	36 861609 36 861609	37 484579	57.016562	25.290338	93.035121	2.3749261	55.785379	4.2591626 	22.1 300/223	95.684196	29.176738	38.526368	91.074661	13.212041	4.9175708	21.112231	0.9735828	75.095947	30.055222	39.4357068	67.238228	25.232533	:64.067883 55 55005	33.952168	17.380996	12.668931	41.987362	9.7453857	06.160722	1.4137855	4.7988078	CI 1.2020.2	ECT 104 159	94.437763	51.377791	37.264497	2.1633354	4.3314462	38.866584 07.135479
	cc cc	135705 5	506582 1	250146 1	1 20029 2 904036 1	17693 3	743313 1	132258 1	999971 1	395857 8	315542 1	736988 9	0 000116	2//0/4 923412 1	396835 3	710527 2	709146 2	972415 1	264795 9	974803 1	917305 6	351/56 1	500000 C	74235 3	316586 1	340044 2	2053307 2	937313 2	520453 1	332848 1	457517 1	310944 9	207321 1	368887 1	084362 1	4100/4 8	1 105010	070907 2	902632 1	159114 2	461795 8	12741 6	(49693 1 133887 1
_ 0000		149 22.24	31 19.4	869 22.33	11 23.0	0.02 07.	57 21.47	52 19.34	56 21.19	155 23.1	66 21.7:	83 20.97 1 8 22 0.97	10 22.01	16.3 1979	163 23.88	91 21.47	49 24.07	21.6	17 22.3	59 21.39	891 20.19	126 20.98	1.22 100	18.2 18.2	35 20.6	871 24.63	82 21.20	125 23.19	33 21.75	242 22.6	504 21.4	324 22.28	801 22.77	73 22.6:	199 21.0	77.07 QL	19.91	126 22.80	55 20.79	041 23.3	24 18.54	11 20.74	77 20.7
	τŰ	10.20460	10.93826	12.68018		14 85367	10.80277	9.714895	12.14487	11.47913	11.53071	8.549706 6 514204	0.2410.0	12 71004	16.9364	14.85358	15.62717	10.40182	9.554523	10.24685	8.754123	7 200505	2000000	7.026198	11.86655	13.9793	15.42551	15.06900	9.76092	9.771612	11.71795	10.30106	9.031833	7.069861	4.553496	10.9390	13 488.20	15.2154	14.30137	15.32536	9.556627	8.867701	10.76725
	<u>ں</u> م	.9786669	.8688859	.9501721	4100747	21 62255	2616718	.8828969	.4257561	.2013736	.8715731	.6484877	. 1900039	6250011	1065833	.1334212	.5412318	.9487906	.8431223	.9718591	5028837	.7864637	.1 230001	.1847366	.2060796	.7897859	.3050891	.5546704	.4642795	.0447937	9.285746	.0504593	.2781653	.1866094	18492011	00000000	0111208.	.2041723	.3915742	.2906278	.9073679	.2803255	.2230828 .1192163
	<u> </u>	1	15	16	10 2	2	1	1 17	1 19	1 18	1 17			1 0	212	22	21	1 18	1 16	1 16	1 4			- 5	18	19	22	53 -0	1 16	1 19	-	1 19	1 16	1 13	1 7.7	<u></u>	<u> </u>	212	18	21	1 16	1 16	1 20
	00						~	-	-	125	-		_					333	-	-	. .			_					-	-	-	1.8	. -	-	-						-	. .	
	00						-	-	-	۔ ب	.		-					1 1.08	-	-	. .		- •	-						-	-	.	. -	.	923						-	. .	
	L C						-	-	-	-	-		_					+	-	-	÷ -			_					-	-	-	.	~ .	-	1 1.076						-	~ .	
	ш ш с						66667	33333	1.3125	-	1.5	14286 6667	10000					33333	84615	33333	45455	1.75		1429					1.625	33333	60606		33333	33333	76923						-	<u>.</u>	
	2 Ш	80	88	5 2	2 2	1 0		72 1.133	51	37	*	28 1.357	014.1	<u> </u>	2 9	2	57	55 1.083	46 1.153	74 1.533	96 1.545	4 0	11 1 20E	11 1.203	37	28	75	9 9	67	72 1.583	36 2.090	36	29 1.133	17 1.333	53 1.230 of	ខ្ល	8 2	8	35	74	28	8	£ 8
$> \vdash \square$	шш	0.064954	0.122428	0.151394:	19708012.0	0.32398	0.15073	0.120278	0.173731	0.148274	0.140206	0.096952	0.0593090	0.187856	0.296259	0.286231	0.279431	0.127363	0.105139	0.116267	0.079819	0.210115	2101000	0.047322	0.16054	0.202709:	0.380257	0.34074:	0.10564:	0.135202	0.185874:	0.143633	0.101914:	0.054786	0.016392	0.101407.0	0 165508	0.264993	0.181653:	0.28471	0.107559:	0.092637:	0.153821
F	0 L	0		0		10 5	0	0	0	0	0	12.5	0		18.75	12.5	12.5	0	0	0	0 0	0 0		0 0			18.75	12.5	6.25	0	0	0	12.5	0	0 0	0 1	0./0	25	0	25	0	0	12.5 12.5
шОнд	οv						3.45	2.52	3.53333333	3.7		0.9	671.7					3.6	2.21428571	1.8	3.3	4.56	0.1	0.34					ę	2.52	1.1	3.2125	3.6	2.35	0.85						4.18571429	3.3	2.88333333
∪ro≩zı	0 F						65.8831209	53.6257016	69.917734	73.8001852	68.7753532	73.6382918	10.2201022.01					80.1826955	57.6331055	75.7636649	63.2802191	66.5894164	4100/10.00	04.1103200					78.3596719	79.0381409	75.9890246	69.4306921	81.7586126	62.9166667	67.047504						75.8043559	73.2017587	/6.144/805 73.2157926
- z ⊢ z o	Ωш						0.94680734	1.34544877	1.24043448	1.36406245	1.55334619	1.35386068 0 0075207	18001070.0					1.22184921	1.08336192	0.9071169	0.97017538	1.04653287	1.0012800.1	1200000000					1.36664646	0.8141809	0.94083136	0.99050333	0.94051552	1.11666667	0.84582334						0.87608532	1.28911616	0.87948832 1.1125845
ш О । – Z О І	ш×						12.0214815	31.0328235	9.3842246	3.45945946		31.7180312 21.2720E02	7600/1717					11.759578	11.2367035	6.76193062	4.74537037	3.25811809	C/00.C7	20.1 333333					36.0520833	11.2049708	36	25.1011048	3.36111111	2.22463768	23.4375						2.27681941	19.1887446	39.9274074
$\square \square \triangleleft \triangleleft \triangleleft \square \triangleleft \sqcap \land \vdash $	- 0		6.25		12.5	2.4	0.21848099	0.30168029	0.16264974	0.1926254	0.19920915	0.24269199	0.20210423	C	>			0.26730986	0.33538093	0.363101	0.2532862	0.19894465	0.2024320	0.33010004	12.5				0.23188741	0.20663287	0.24781772	0.2155214	0.22669213	0.27875254	0.36508825						0.26542015	0.22594353	0.15724924
ω — ш :	Σთ	0	0	60606060.6	10	7 14285714	0	6.66666667	0	12.5	37.5	0 0	D	C	28.5714286	0	6.66666667	8.333333333	7.69230769	0	0	8.3333333333	U 7 1 4 7 2 5 7 1 4	41/2027/17/16.6666667	0		11.1111111 25	0 7	12.5	8.333333333	0	0	13.3333333	0	0	30.3030304	Ð	0	66.6666667	16.6666667	0	0	00
ω — ш	∑∞	30.7692308	0	0	10	7 14285714	6.6666667	0	18.75	0	6.25	7.14285714	0.000000000	C	28.5714286	8.333333333	33.3333333	8.33333333	7.69230769	0	0 0	0 0	14 2867142	16.6666667	0		22.222222	27.2727233	6.25	8.333333333	9.09090909	0	6.66666667	0	7.69230769	0	41.142821.1	15.3846154	0	16.6666667	18.1818182	0	0.05290769
ω – ш	×⊼	30.7692308	0	36.3636364	40	57 1428571	33.33333333	13.3333333	18.75	0	18.75	14.2857143 16 666667	1000000101	C	7.14285714	33.3333333	13.3333333	25	30.7692308	7.14285714	45.4545455	50 12 E	7 14726714	1. 14203/ 14	0		22.222222	27.272733	43.75	33.3333333	18.1818182	30	20	0	30.7692308 54 545 45 45	0404040.40	D	15.3846154	0	25	18.1818182	50	15.3846154 40
о ー ш	Σo	23.0769231	9.09090909	36.3636364	20	21 4285714	46.6666667	73.3333333	37.5	62.5	31.25	21.4285714 E0	ne	6 6666667	28.5714286	25	13.3333333	25	38.4615385	21.4285714	36.3636364	8.333333333 17 E	12.21 1 1 1225714	41.6666667	28.5714286		44.444444	27.27273 27.27273	37.5	33.3333333	36.3636364	40	40	66.6666667	15.3846154		41/02/14/	38.4615385	33.3333333	8.33333333	36.3636364	20	23.0769231
о ー ш	o Z	15.3846154	0	9.09090909	10	7 14285714	0	6.66666667	12.5	0	6.25	28.5714286	67	06	3 0	25	26.6666667	33.3333333	7.69230769	14.2857143	9.090909090 20	25	02 024730	6741700.24 0	21.4285714			0.33333333	0	16.6666667	0	20	20	0	30.7692308	000772200	28.51/12/20	7.69230769		8.33333333	18.1818182	20	1.69230769
ω н ш	Σ4	0	0 0	0 24 25	31.25 10		6.66666667	0	6.25	0	0	21.4285714		U 46 6666667	7.14285714	0	6.66666667	0	7.69230769	21.4285714	60606060.6	8.333333333 27 E	0.10	0 0	28.5714286	0	c	0 0	0	0	9.09090909	0	0	33.33333333	15.3846154	9.09090909	0074170.07	23.0769231			0	10	38.4613385
о ⊢ ш :	Σ ო	0	36.3636364	9.090909090 0		• c	0	0	6.25	25	0	7.14285714		26 666667	0	8.33333333	0	0	0	21.4285714	0 0	0 u	7 14796714	41/00241./	7.14285714	0	c	0 0	0	0	0	10	0	0	0 0	0	41.142821.1	0			0	0	00
о — Ш :	ΣN	0	45.454545 2	0 0			0	0	0	0	0	00		0.7L	0		0	0	0	7.1428571	0 0	0 0		0 0	14.285714	6.25	c	0 0	0	0	9.0909091	0	0	0	0 0	0 12200 11	41/C07.41	0			9.0909091	0	0

Plot Data Set - Habinsaran